


Concentrations of *Escherichia Coli* in Streams in the Upper Wabash River Watershed in Indiana, June–September 1998

Water-Resources Investigations Report 00–4021



*Prepared in cooperation with the
Indiana Department of Environmental Management*

U.S. Department of the Interior
U.S. Geological Survey

Concentrations of *Escherichia Coli* in Streams in the Upper Wabash River Watershed in Indiana, June–September 1998

By Cheryl A. Silcox, David C. Voelker, and Timothy C. Willoughby

Water-Resources Investigations Report 00-4021

Prepared in cooperation with the

Indiana Department of Environmental Management

Indianapolis, Indiana

2000

U.S. Department of the Interior
BRUCE BABBITT, Secretary

U.S. Geological Survey
Charles G. Groat, Director

The use of trade, product, or firm names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

For additional information, write to:

District Chief
U.S. Geological Survey
Water Resources Division
5957 Lakeside Boulevard
Indianapolis, IN 46278-1996

Copies of this report can be purchased from:

U.S. Geological Survey
Branch of Information Services
Box 25286
Federal Center
Denver, CO 80225-0286

Contents

| | |
|--|----|
| Abstract | 1 |
| Introduction | 1 |
| Purpose and Scope | 1 |
| Description of the Study Area | 2 |
| Selection of Sampling Sites | 2 |
| Methods of Sampling | 2 |
| Field Measurements | 2 |
| Collection of Samples | 7 |
| Processing of Samples | 7 |
| Concentrations of <i>Escherichia coli</i> | 8 |
| Quality-Assurance and Quality-Control Procedures | 25 |
| Filter Blanks | 27 |
| Process Blanks | 27 |
| Field Blanks | 27 |
| Summary | 27 |
| References Cited | 28 |
| Supplemental Data | 29 |

Figures

1–4. Maps showing:

1. Location and principal streams of the Upper Wabash River Watershed in Indiana 3
2. Location of *Escherichia coli* sampling sites within the Upper Wabash River Watershed 4
3. Ranges in concentrations of *Escherichia coli* for each site 9
4. Five-sample geometric mean *Escherichia coli* concentration computed for each site10

5–7. Graphs showing:

5. Concentrations of *Escherichia coli* and five-sample geometric means for selected sites12
6. Stream discharge and concentrations of *Escherichia coli* at selected sites16
7. Concentrations of *Escherichia coli* measured in the environmental samples and duplicate samples and their natural log percent difference26

Contents—Continued

Tables

| | |
|--|----|
| 1. Sites at which water samples were collected during June and July 1998 (Group 1 sites) for analysis of <i>Escherichia coli</i> , Upper Wabash River Watershed in Indiana | 5 |
| 2. Sites at which water samples were collected during August and September 1998 (Group 2 sites) for analysis of <i>Escherichia coli</i> | 6 |
| 3. Water-quality data for selected sites | 30 |
| 4. Water-quality data for two sites in the Upper Wabash River Watershed in Indiana where additional samples were collected during July–September 1998. | 42 |
| 5. Quality-assurance data associated with the Group 1 <i>Escherichia coli</i> samples | 43 |
| 6. Quality-assurance data associated with the Group 2 <i>Escherichia coli</i> samples | 48 |

Conversion Factors and Abbreviations

| Multiply | By | To obtain |
|---|---------|-----------------------------|
| square mile (mi ²) | 2.590 | square kilometer |
| cubic foot per second (ft ³ /sec) | 0.02832 | cubic meter per second |
| pounds per square inch (lbs/in ²) | 70.307 | grams per square centimeter |

Temperature is given in degrees Celsius (°C), which can be converted to degrees Fahrenheit (°F) by use of the following equation:

$$^{\circ}\text{F} = 1.8(^{\circ}\text{C}) + 32$$

Abbreviated water-quality units used in this report: Chemical concentrations and water temperature are given in metric units. Chemical concentration is given in milligrams per liter (mg/L) or micrograms per liter (µg/L). Milligrams per liter is a unit expressing the concentration of chemical constituents in solution as weight (milligrams) of solute per unit volume (liter) of water. One thousand micrograms per liter is equivalent to one milligram per liter. For concentrations less than 7,000 mg/L, the numerical value is the same as for concentrations in parts per million. Concentrations of bacteria are given in colonies per 100 milliliters (col/100 mL).

Specific conductance of water is expressed in microsiemens per centimeter at 25 degrees Celsius (µS/cm). This unit is equivalent to micromhos per centimeter at 25 degrees Celsius (µmho/cm), formerly used by the U.S. Geological Survey.

Volumes of water-quality samples are given in liters (L) and milliliters (mL).

Concentrations of *Escherichia Coli* in Streams in the Upper Wabash River Watershed in Indiana, June–September 1998

By Cheryl A. Silcox, David C. Voelker, and Timothy C. Willoughby

Abstract

Water samples collected from 46 stream sites in the Upper Wabash River Watershed from June through September 1998 were analyzed for concentrations of the bacteria *Escherichia coli* (*E. coli*). Each site was sampled five times in a 30-day period. Twenty-one sites were sampled during June and July, and 25 sites were sampled during August and September. The concentration of *E. coli* in 145 of the 230 samples collected exceeded the Indiana standard of 235 colonies per 100 milliliters for a single sample for waters used for recreation. A five-sample geometric mean also was computed for each site. Concentrations in samples from 43 of the 46 sites exceeded the Indiana bacteriological quality standard of 125 colonies per 100 milliliters for a five-sample geometric mean for waters used for recreation.

Discharge during sample collection generally was greater than the long-term median daily mean discharge. To determine if the greater discharge affected the concentrations of *E. coli*, additional samples were collected at two sites. Statistically significant positive correlations between concentrations of *E. coli* and discharge were determined for these sites, indicating increased concentrations with greater discharge.

Introduction

The Indiana Department of Environmental Management (IDEM) is responsible for monitoring the quality of Indiana's waters. IDEM monitors watersheds on a rotating basis and selected the Upper Wabash River Watershed for monitoring during 1998. As part of this effort, IDEM entered into a cooperative program with the U.S. Geological Survey (USGS) to measure concentrations of the bacteria *Escherichia coli* (*E. coli*) at 46 stream sites in the Upper Wabash River Watershed from June through September 1998.

The presence of *E. coli* in water is direct evidence of the presence of fecal contamination from warm-blooded animals and indicates the possible presence of pathogens (Myers and Sylvester, 1997). *E. coli* is one of the two preferred indicator bacteria used by the U.S. Environmental Protection Agency (USEPA) to determine the suitability of surface waters for recreational use. The water-quality standards for *E. coli* in recreational waters in Indiana require the concentration of *E. coli* to be less than a single-sample standard or less than the geometric mean computed from five samples collected within a 30-day period (Oddi, 1995).

Purpose and Scope

This report documents the concentrations of *E. coli* measured in samples from selected streams in the Upper Wabash River Watershed from June through September 1998. The report describes the relation between concentrations of *E. coli* and streamflow at sites where streamflow records were available and presents the quality-assurance data

for the *E. coli* samples. Field measurements of water temperature, pH, dissolved oxygen, specific conductance, and turbidity also are presented.

Description of the Study Area

The part of the Upper Wabash River Watershed that was sampled extends from the Indiana-Ohio state line downstream to include Wildcat Creek near Lafayette, Ind. (fig. 1). The watershed drains 6,918 mi² in Indiana and 285 mi² in Ohio, for a total of 7,203 mi² (Hoggatt, 1975). The predominant land use in the watershed is agricultural (row crop and pasture).

The Upper Wabash River Watershed lies within two distinct physiographic areas (Schneider, 1996)—the Northern Moraine and Lake Region, located north of the Eel River, and the Tipton Till Plain, located predominantly south of the Eel River. The land surface generally is flat to gently undulating, poorly drained, and featureless.

Three principal reservoirs in the basin were built as flood-control structures and have a dual role as recreational water bodies. They are Huntington, Salamonie, and Mississinewa Lakes. Major tributaries to the Wabash River in the study area include the Little, Salamonie, Mississinewa, Eel, and Tiptecanoe Rivers, and Pipe, Deer, and Wildcat Creeks.

Selection of Sampling Sites

Sampling-site locations initially were selected by IDEM personnel. Responses from a 1987 poll of local health officials, conservation officers, and sheriff's departments regarding known areas of stream recreational uses were used to provide a core list of potential sampling sites. Additional sites then were added for spatial coverage and (or) positioning of sites at existing USGS streamflow-gaging stations. Site locations were plotted on topographic maps and verified in the field by USGS personnel prior to initiating sample collection during June 1998. Where sampling conditions were unsafe or where site characteristics interfered with the samplers' ability to collect a sample, the site

was relocated as close to the initial site as possible. Changes to sampling-site locations were agreed upon by USGS and IDEM personnel.

The 46 sampling sites selected were divided into two groups so that all sites could be sampled five times at equally spaced intervals within a 30-day period. Eight additional samples were collected weekly from July 27 to September 9, 1998, at sites 2 and 15. Figure 2 shows the locations of the sampling sites. Table 1 lists the Group 1 sites (sites 1 through 21) that were sampled during June and July 1998. Table 2 lists the Group 2 sites (sites 22 through 46) that were sampled during August and September 1998.

Methods of Sampling

Water samples were collected during the recreational season in Indiana, defined as April through October. The samples usually were collected by two-person field crews to expedite the sampling process and meet the mandated 6-hour sample-holding-time limit prior to processing the samples. Duties at the sampling sites included completing the field forms, measuring and recording water-surface-elevation data, documenting site characteristics at the time of sampling, measuring and recording field parameters, inspecting the streamflow-gaging station if one were present, and collecting the samples.

Field Measurements

A multi-parameter water-quality measuring meter was used to make field measurements of water temperature, dissolved oxygen, pH, and specific conductance at several locations across the width of the stream. The meter was calibrated daily for dissolved oxygen, pH, and specific conductance before any field measurements were made. Field determinations of turbidity were made by collecting samples of surface water in polyethylene bottles and analyzing the samples with a portable turbidimeter. The measuring range of the portable turbidimeter was checked daily with reference standards. If parameters measured in the field did





Figure 1. Location and principal streams of the Upper Wabash River Watershed in Indiana.



EXPLANATION

- WATERSHED BOUNDARY
- COUNTY BOUNDARY
- GROUP 1 *ESCHERICHIA COLI* SAMPLING SITES --
Sites sampled June and July 1998
- GROUP 2 *ESCHERICHIA COLI* SAMPLING SITES--
Sites sampled August and September 1998

Figure 2. Location of *Escherichia coli* sampling sites within the Upper Wabash River Watershed in Indiana.

Table 1. Sites at which water samples were collected during June and July 1998 (Group 1 sites) for analysis of *Escherichia coli*, Upper Wabash River Watershed in Indiana

[USGS, U.S. Geological Survey; SR, State Road; CR, County Road; E, W, N, and S denote the geographic directions of east, west, north, and south]

| Site number | Site location | USGS site identification | Latitude and longitude | | County |
|-------------|---|--------------------------|------------------------|------------|------------|
| 1 | Wabash River at State Line Road | 03322500 | 40°33' 50" | 84°48' 10" | Jay |
| 2 | Wabash River at SR 218 at Linn Grove | 03322900 | 40°39' 22" | 85°01' 58" | Adams |
| 3 | Wabash River at CR 300N near Bluffton | 404709085120001 | 40°47' 09" | 85°12' 00" | Wells |
| 4 | Little River at CR 200E near Huntington | 405354085244701 | 40°53' 54" | 85°24' 47" | Huntington |
| 5 | Wabash River at SR 105 near Andrews | 405208085360601 | 40°52' 08" | 85°36' 06" | Huntington |
| 6 | Wabash River at Wabash | 03325000 | 40°47' 25" | 85°49' 13" | Wabash |
| 7 | Salamonie River at CR 75S near Portland | 03324200 | 40°25' 40" | 85°02' 20" | Jay |
| 8 | Salamonie River at SR 18 at Matamoras | 403316085154201 | 40°33' 16" | 85°15' 42" | Blackford |
| 9 | Salamonie River at SR 124 near Lancaster | 404430085303201 | 40°44' 30" | 85°30' 32" | Huntington |
| 10 | Wabash River at Broadway Street at Peru | 404432086054801 | 40°44' 32" | 86°05' 48" | Miami |
| 11 | Pipe Creek at CR 125W near Bunker Hill | 03327520 | 40°40' 06" | 86°05' 44" | Miami |
| 12 | Mississinewa River at CR 100W near Ridgeville | 03325500 | 40°16' 48" | 84°59' 33" | Randolph |
| 13 | Mississinewa River near Walnut Road near Eaton | 402026085221001 | 40°20' 26" | 85°22' 10" | Delaware |
| 14 | Mississinewa River at CR 950E near Matthews | 402339085293601 | 40°23' 39" | 85°29' 36" | Grant |
| 15 | Mississinewa River at Highland Avenue at Marion | 03326500 | 40°34' 34" | 85°39' 34" | Grant |
| 16 | Mississinewa River at CR 500N at Jalapa | 403732085435601 | 40°37' 32" | 85°43' 56" | Grant |
| 17 | Eel River at SR 9 near Columbia City | 410733085285201 | 41°07' 33" | 85°28' 52" | Whitley |
| 18 | Eel River at SR 14 at North Manchester | 03328000 | 40°59' 55" | 85°45' 50" | Wabash |
| 19 | Eel River at SR 16 near Roann | 405448085563901 | 40°54' 48" | 85°56' 39" | Miami |
| 20 | Eel River at Meridian Road near Denver | 405125086043301 | 40°51' 25" | 86°04' 33" | Miami |
| 21 | Eel River at Adamsboro Road near Logansport | 03328500 | 40°46' 55" | 86°15' 50" | Cass |

Table 2. Sites at which water samples were collected during August and September 1998 (Group 2 sites) for analysis of *Escherichia coli*, Upper Wabash River Watershed in Indiana

[USGS, U.S. Geological Survey; SR, State Road; CR, County Road; E, W, and N denote the geographic directions of east, west, and north]

| Site number | Site location | USGS site identification | Latitude and longitude | | County |
|-------------|--|--------------------------|------------------------|------------|------------|
| 22 | Tippecanoe River at SR 5 at Wilmot | 411832085383901 | 41°18' 32" | 85°38' 39" | Noble |
| 23 | Tippecanoe River at Oswego | 03330500 | 41°19' 14" | 85°47' 21" | Kosciusko |
| 24 | Tippecanoe River at CR 700W near Atwood | 411438085462001 | 41°14' 38" | 85°46' 20" | Kosciusko |
| 25 | Tippecanoe River at SR 331 near Old Tip Town | 411308086065401 | 41°13' 08" | 86°06' 54" | Marshall |
| 26 | Tippecanoe River at Old US 31 near Rochester | 410620086131301 | 41°06' 20" | 86°13' 13" | Fulton |
| 27 | Tippecanoe River at Leiter's Ford | 410730086225901 | 41°07' 30" | 86°22' 59" | Fulton |
| 28 | Tippecanoe River at CR 200E near Ora | 03331500 | 41°09' 26" | 86°33' 49" | Pulaski |
| 29 | Tippecanoe River at SR 119 near Winamac | 410023086361001 | 41°00' 23" | 86°36' 10" | Pulaski |
| 30 | Tippecanoe River at Buffalo | 03332345 | 40°53' 05" | 86°44' 49" | White |
| 31 | Big Monon Ditch at SR 16 near Buffalo | 405208086464401 | 40°52' 08" | 86°46' 44" | White |
| 32 | Tippecanoe River at SR 24 at Monticello | 404446086452701 | 40°44' 46" | 86°45' 27" | White |
| 33 | Tippecanoe River at SR 18 near Delphi | 03333050 | 40°35' 38" | 86°46' 12" | Carroll |
| 34 | Middle Fork Wildcat Creek at CR 775E | 402505086452301 | 40°25' 05" | 86°45' 23" | Tippecanoe |
| 35 | South Fork Wildcat Creek at SR 26 near Lafayette | 03334500 | 40°25' 04" | 86°46' 05" | Tippecanoe |
| 36 | Wildcat Creek at CR 2A near Lafayette | 03335000 | 40°26' 26" | 86°49' 45" | Tippecanoe |
| 37 | Wabash River at SR 225 near Battleground | 402943086492301 | 40°29' 43" | 86°49' 23" | Tippecanoe |
| 38 | Wabash River at CR 200N near Delphi | 403448086413401 | 40°34' 48" | 86°41' 34" | Carroll |
| 39 | Wabash River at CR 675W near Georgetown | 404420086301001 | 40°44' 20" | 86°30' 10" | Cass |
| 40 | Wildcat Creek at CR 1100E near Jerome | 03333450 | 40°26' 29" | 85°55' 08" | Howard |
| 41 | Wildcat Creek at SR 22 near Burlington | 402920086221201 | 40°29' 20" | 86°22' 12" | Howard |
| 42 | Deer Creek at SR 29 at Deer Creek | 403630086232801 | 40°36' 30" | 86°23' 28" | Carroll |
| 43 | Deer Creek at CR 300N near Delphi | 03329700 | 40°35' 25" | 86°37' 15" | Carroll |
| 44 | Wildcat Creek at SR 39 at Owasco | 03334000 | 40°27' 50" | 86°38' 15" | Carroll |
| 45 | Kilmore Creek at CR 600W near Hamilton | 402009086370001 | 40°20' 09" | 86°37' 00" | Clinton |
| 46 | South Fork Wildcat Creek near Hamilton | 03334230 | 40°19' 15" | 86°37' 06" | Clinton |

not appear stable or have reasonable values, the measuring meter was recalibrated at the site and field parameters were remeasured.

Collection of Samples

Water samples for *E. coli* determinations were collected in 300-mL (milliliter) glass bottles with glass stoppers. Prior to use, the bottles were washed with detergent, rinsed three times with hot water and three times with deionized water, and sterilized by autoclaving. To ensure optimum growing conditions for *E. coli*, two solutions were added to each bacteria-sample bottle before the bottle was sterilized. A 10-percent solution of sodium thiosulfate was added to counter the effects of residual chlorine or other halogens used in water-disinfection processes. Residual chlorine and other halogen compounds act as bacterial-growth inhibitors; their effects need to be reduced so that the *E. coli* can be recovered fully on growth medium and produce accurate counts (Bordner and Winter, 1978; American Public Health Association and others, 1992). In addition, a 15-percent solution of ethylenediaminetetraacetic acid (EDTA) was added to neutralize the effects of trace-element concentrations greater than 10 µg/L (micrograms per liter). EDTA, a chelating agent, binds particularly with copper and zinc, making the metals neutral so that they do not adversely affect bacterial growth (Britton and Greeson, 1989, p. 5–6; Bordner and Winter, 1978; American Public Health Association and others, 1992).

In the field, a weighted hand-line sampler, which held the sample bottle, was lowered beneath the surface of the water in the center of flow. Some samples were collected by immersing the bottles by hand when the stream was too shallow for the hand-line sampler. At each site, a sample was a composite of water from one to six well-mixed areas of flow, depending on the width of the stream. The samples were kept chilled until processing.

Processing of Samples

Equipment used to process the samples was washed with detergent prior to field work, rinsed

three times with tap water and three times with deionized water, and then sterilized with an 8-watt ultraviolet (UV) lamp with a wave length of 254 nanometers for a minimum of 15 minutes. Processing equipment included a UV sterilization chamber, a multi-port manifold filter stand, stainless-steel filter holders, pumps, sterile disposable pipets, glass graduated cylinders, refrigeration units, and incubators. At least two aluminum-block incubators capable of maintaining temperatures of 35.0°C and 44.5°C were used to provide optimum conditions for bacterial growth.

The *E. coli* substrate medium kit, prepared by the USGS Quality of Water Services Unit (QWSU) in Ocala, Fla., according to USEPA Method 1103.1, (Bordner and others, eds., 1978) was used to prepare the membrane-filter thermotolerant *E. coli* media (mTec) agar. The mTec agar was prepared in the USGS Indiana District laboratory for use in the field. The 2-week holding time for mTec agar, once it was prepared, was monitored in the laboratory and field. Urea/Phenol Red reagent, also included in the kit, was prepared in the field and used to confirm the presence of *E. coli* colonies.

The samples collected for analysis of *E. coli* were processed either in the field in hotel rooms or in the USGS laboratory in Indianapolis. Surfaces on which the samples were processed were cleaned with isopropyl alcohol before the first sample was processed each day, between samples, and after the last sample was processed each day. Sample handlers washed their hands with bactericidal soap before processing the first sample, between samples, and after processing the last sample.

Sample dilutions were made by adding 11 mL of sample water to 99 mL of sterile dilution water for a 1:10 ratio and 1 mL of sample water to another 99 mL of sterile dilution water for a 1:100 dilution. Five to eight different sample volumes, including one to three different dilutions, were filtered for each site to obtain at least one sample volume that represented the ideal colony count of 20 to 80 colonies per filter plate (Myers and Sylvester, 1997).

Sterile, disposable 1- and 10-mL glass pipets were used to measure and deliver concentrated sample volumes to dilution bottles and to measure and deliver dilution volumes to the interior of the funnel filter assembly. For sample dilution volumes less than 10 mL, about 20 mL of sterile saline buffer solution was poured into the funnel before pipetting the sample dilution to evenly distribute the bacteria on the filter. A sterile graduated cylinder was used to transfer sample dilution volumes greater than 10 mL. Samples were shaken vigorously before each sample dilution volume was withdrawn. A three-port manifold with funnels or a single-use stainless-steel filter system was used to support a 0.45- μm (micrometer) filter. The water was pulled through the filter either by a vacuum pump set not to exceed 5 lb/in² (pounds per square inch) of pressure or by a hand-vacuum assembly. After filtering each of the sample dilution volumes, 20 to 30 mL of sterile saline buffer solution were used to flush the sides of the funnel to ensure that any bacteria present on the funnel walls were rinsed onto the filter. The graduated cylinders used to measure and deliver sample dilution volumes to the funnel also were rinsed with sterile saline buffer solution, and the rinsate was processed through the filter.

Petri dishes were labeled prior to processing the sample. Concentrated sample and sample dilutions were filtered from smallest to largest. The filters then were placed in petri dishes with the mTec agar and placed inverted in a pre-heated incubator set at 35.0°C for 1.75 to 2 hours, removed, and then placed in a pre-heated incubator set at 44.5°C for 22 to 24 hours. After the second incubation period was completed, the filter was transferred to a filter pad saturated with Urea/Phenol Red reagent. After 15 to 20 minutes at room temperature, only the yellow to yellow-brown *E. coli* colonies were counted. If the filter plate had a colony count in the ideal range, verification of the count was made either by the second crew member or by rotating the filter 90 degrees and the same crew member recounting the colonies. If more than one dilution were within the ideal colony count, the concentration of *E. coli* was computed as the sum of the colony counts for each sample volume, multiplied by 100, and divided by the sum of the dilution


volumes. The same calculation was used if none of the dilutions had concentrations of *E. coli* within the ideal colony count, and the result is reported as an estimate. (A “K” before the value indicates that the values are estimated from non-ideal colony counts.) Concentrations of *E. coli* were calculated according to the methods described by Myers and Sylvester (1997) and recorded on the field sheet for each site. Concentrations of *E. coli* are reported in whole numbers for results less than 10, and two significant figures are reported for results greater than or equal to 10 (Myers and Wilde, 1997). After being counted, *E. coli* petri dishes were sterilized with chlorine bleach, placed in sealed plastic bags, and disposed.

Concentrations of *Escherichia Coli*

The Indiana environmental rules establish the bacteriological quality standard for waters for recreational uses (Oddi, 1995, 327 IAC 2-1-6 [d]). These rules are used to evaluate waters for full-body-contact recreational uses, to establish wastewater-treatment requirements, and to establish effluent limits during the recreational season, which is defined as the months of April through October. The standard states that:

E. coli bacteria, using membrane filter (MF) count, shall not exceed one hundred twenty-five (125) per one hundred (100) milliliters as a geometric mean based on not less than five (5) samples equally spaced over a thirty (30) day period nor exceed two hundred thirty-five (235) per one hundred (100) milliliters in any one (1) sample in a thirty (30) day period.


Table 3, at the back of the report, lists field measurements and concentrations of *E. coli* for all 46 sites. The five-sample geometric mean is shown below the last collection date for each site. Figure 3 shows the range of concentrations of *E. coli* for the five samples collected at each site, and figure 4 shows the five-sample geometric-mean concentration computed for each site.



EXPLANATION

-  WATERSHED BOUNDARY
-  COUNTY BOUNDARY
-  130
GROUP 1 *ESCHERICHIA COLI* SAMPLING SITES--
Showing range of colonies per 100 milliliters
-  100
GROUP 2 *ESCHERICHIA COLI* SAMPLING SITES--
Showing range of colonies per 100 milliliters
- K** Values estimated from non-ideal colony counts

Figure 3. Ranges in concentrations of *Escherichia coli* for each site in the Upper Wabash River Watershed in Indiana, June-September 1998.



EXPLANATION

- BASIN BOUNDARY
- COUNTY BOUNDARY
- GROUP 1 *ESCHERICHIA COLI* SAMPLING SITES--
Showing five-sample geometric mean per 100 milliliters
- GROUP 2 *ESCHERICHIA COLI* SAMPLING SITES--
Showing five-sample geometric mean per 100 milliliters

Figure 4. Five-sample geometric mean *Escherichia coli* concentration computed for each site in the Upper Wabash River Watershed in Indiana, June-September 1998.

The five-sample geometric mean of concentrations of *E. coli* for all sites ranged from 17 to 4,800 colonies per 100 mL, and the concentrations in samples from 43 sites exceeded the five-sample geometric-mean standard. Concentrations of *E. coli* at all 46 sites ranged from an estimated 6 to 110,000 colonies per 100 mL, with concentrations in 145 of the 230 samples processed exceeding the single-sample standard. At the three sites where concentrations did not exceed the five-sample geometric-mean standard, only one sample had a concentration that exceeded the single-sample standard.

Figure 5 shows the concentration of *E. coli* at sampling sites in the watershed. Sites are grouped by location along the Wabash, Salamonie, Mississinewa, Eel, and Tiptecanoe Rivers, Wildcat Creek, and by other tributaries. Concentrations of *E. coli* at all nine sites along the Wabash River exceeded the standard for the five-sample geometric mean, and concentrations for 28 of the 45 samples processed from those sites exceeded the standard for a single sample. Concentrations of *E. coli* at the Wabash River sites ranged from 20 to 110,000 colonies per 100 mL, and the five-sample geometric mean ranged from 150 to 3,100 colonies per 100 mL.

Thirty-five samples from eight sites on the Salamonie and Mississinewa Rivers had concentrations of *E. coli* that exceeded the single-sample standard. Concentrations of *E. coli* in samples from sites on the Salamonie River ranged from 170 to 49,000 colonies per 100 mL and from 130 to 19,000 colonies per 100 mL in samples from sites on the Mississinewa River. Concentrations at all sites on the Salamonie and Mississinewa Rivers exceeded the standard for the five-sample geometric mean. For the Eel River, 23 samples from five sites had concentrations that exceeded the single-sample standard, and concentrations at all sites exceeded the five-sample geometric-mean standard. Concentrations of *E. coli* in samples from the Eel River ranged from 48 to 17,000 colonies per 100 mL, and the five-sample geometric mean ranged from 490 to 1,400 colonies per 100 mL.

At 8 of the 11 sites on the Tiptecanoe River, concentrations of *E. coli* exceeded the standard for

the five-sample geometric mean. Concentrations of *E. coli* in samples from the Tiptecanoe River sites ranged from an estimated 6 to 70,000 colonies per 100 mL, and the five-sample geometric mean ranged from 17 to 670 colonies per 100 mL. Only sites 22, 23, and 33 had five-sample geometric-mean concentrations that did not exceed the standard. No samples from sites 22 and 23 had concentrations that exceeded the standard for a single sample, and site 33 had one sample that exceeded the standard.

Stream discharges presented in this report were taken from data collected at USGS streamflow-gaging stations located at or near the sampling sites. Figure 6 shows the relation of concentrations of *E. coli* to discharge for sampling sites in the Upper Wabash River Watershed where stage-discharge relations have been developed. During sampling of the Group 1 sites, most of the samples were collected at discharges greater than the long-term median daily mean discharge (Stewart and others, 1999). Seventeen of the 46 sites were at USGS streamflow-gaging stations. Based on records of streamflow from these stations, 62 percent of the samples collected at these sites were collected at discharges above the long-term median daily mean discharge (Stewart and others, 1999). Group 1 sites included nine streamflow-gaging stations, at which 69 percent of the samples were collected at discharges above the long-term median daily mean discharge. Group 2 sites included eight streamflow-gaging stations, at which 55 percent of the samples were collected at discharges above the long-term median daily mean discharge.

Eight additional samples were collected at sites 2 and 15 from July 22 to September 9, 1998, to examine further the effect of variations in flow conditions on the concentrations of *E. coli*. Figure 6 and table 4, at the back of this report, display the results for these two sites. A Kendall's Tau test for significant correlation indicated that at both sites, the correlation between discharge and concentration of *E. coli* was statistically significant. For both sites, the slope was positive, indicating increased concentrations of *E. coli* with greater discharge. The plots for sites 2 and 15 (fig. 6) list the probability of obtaining a correlation coefficient between




Figure 5. Concentrations of *Escherichia coli* (*E. coli*) and five-sample geometric means for selected sites in the Upper Wabash River Watershed in Indiana, June–September 1998.




Figure 5. Concentrations of *Escherichia coli* (*E. coli*) and five-sample geometric means for selected sites in the Upper Wabash River Watershed in Indiana, June–September 1998—Continued.




Figure 5. Concentrations of *Escherichia coli* (*E. coli*) and five-sample geometric means for selected sites in the Upper Wabash River Watershed in Indiana, June–September 1998—Continued.

CONCENTRATION OF ESCHERICHIA COLI, IN COLONIES PER 100 MILLILITERS




Figure 5. Concentrations of *Escherichia coli* (*E. coli*) and five-sample geometric means for selected sites in the Upper Wabash River Watershed in Indiana, June–September 1998—Continued.




Figure 6. Stream discharge and concentrations of *Escherichia coli* at selected sites in the Upper Wabash River Watershed in Indiana, June–September 1998.




Figure 6. Stream discharge and concentrations of *Escherichia coli* at selected sites in the Upper Wabash River Watershed in Indiana, June–September 1998—Continued.





Figure 6. Stream discharge and concentrations of *Escherichia coli* at selected sites in the Upper Wabash River Watershed in Indiana, June–September 1998—Continued.





Figure 6. Stream discharge and concentrations of *Escherichia coli* at selected sites in the Upper Wabash River Watershed in Indiana, June–September 1998—Continued.





Figure 6. Stream discharge and concentrations of *Escherichia coli* at selected sites in the Upper Wabash River Watershed in Indiana, June–September 1998—Continued.




Figure 6. Stream discharge and concentrations of *Escherichia coli* at selected sites in the Upper Wabash River Watershed in Indiana, June–September 1998—Continued.




Figure 6. Stream discharge and concentrations of *Escherichia coli* at selected sites in the Upper Wabash River Watershed in Indiana, June–September 1998—Continued.




Figure 6. Stream discharge and concentrations of *Escherichia coli* at selected sites in the Upper Wabash River Watershed in Indiana, June–September 1998—Continued.




Figure 6. Stream discharge and concentrations of *Escherichia coli* at selected sites in the Upper Wabash River Watershed in Indiana, June–September 1998—Continued.

concentrations of *E. coli* and discharge equal to or greater than the computed correlation coefficient.

Quality-Assurance and Quality-Control Procedures

Quality-assurance and quality-control procedures were followed for collection and processing of the samples. These procedures include frequent checking and calibration of equipment as well as collection of additional samples for quality control. Analysis of the quality-control samples provides information not only about the potential for sample contamination during processing but also about the variability of sampling.

The pH buffers and specific-conductance solutions used to calibrate the multi-parameter measuring instrument were quality assured by the QWSU. *E. coli* substrate media kits, membrane filters, sterile saline buffer solution, pre-measured sterile dilution water, petri dishes, and petri dishes with pads also were quality assured by the QWSU. The incubators were checked weekly with an American Society for Testing and Materials (ASTM) certified thermometer to assure that temperature ranges shown on the internal thermometer in the incubator were accurate to $\pm 0.5^{\circ}\text{C}$. The incubators were inspected daily to ensure that they were operating properly.

Quality-control samples consisted of 9 field blanks, 226 filter blanks, 42 process blanks, and 44 duplicate samples. Results of the quality-assurance and *E. coli* determinations are presented in tables 5 and 6 at the back of this report.

During the sampling of the Group 1 sites, 20 sequential duplicate samples were collected at selected sites immediately after the environmental *E. coli* samples were collected. This was accomplished by taking an additional sample from the same location at which the environmental sample was collected. The sequential duplicate samples were processed in the same manner as the environmental samples. The comparison between the environmental samples and the sequential duplicate samples provides information on sampling variability because the sequential duplicate samples were

not collected from exactly the same water as the environmental samples.

Figure 7 displays the difference between the concentrations of *E. coli* measured in the environmental sample and the sequential duplicate samples and their natural log percent difference. The median difference between the environmental samples and the sequential duplicate samples for the Group 1 sites was -10 colonies per 100 milliliters, with a median natural log difference of -3.8 percent. A Wilcoxon signed-rank test (Helsel and Hirsch, 1992) was used to determine if there were any statistically significant differences between the environmental samples and the sequential duplicates. No statistically significant differences were determined between the environmental samples and the sequential duplicate samples for the Group 1 sites at the 5-percent significance level.

During the sampling of the Group 2 sites in August and September, the sequential-duplicate-sampling method used during sampling of the Group 1 sites was replaced with a sampling method that collected concurrent or simultaneous duplicates. Twenty-one concurrent duplicate samples were collected at selected sites in the same location where the environmental samples were collected. The concurrent duplicates were processed in the same manner as the environmental samples and can be used to evaluate the variability in sampling, sampling equipment, and the natural variability in the samples. Figure 7 displays the differences between the concentrations of *E. coli* measured in the environmental samples and the concurrent duplicate samples and their natural log percent difference. The median difference between the environmental samples and the concurrent duplicate samples for the Group 2 sites was 3 colonies per 100 milliliters, with a median natural log difference of 4.7 percent. A Wilcoxon signed-rank test (Helsel and Hirsch, 1992) was used to determine if there were statistically significant differences between the environmental samples and the concurrent duplicates. No statistically significant differences were determined between the environmental samples and the concurrent duplicate samples for the Group 2 sites at the 5-percent significance level.




Figure 7. Concentrations of *Escherichia coli* measured in the environmental samples and duplicate samples and their natural log percent difference.

The similarity between the median natural log percent differences determined for the sequential duplicate samples and the concurrent duplicate samples indicates that the waters sampled generally maintained a constant concentration of *E. coli* for at least short periods of time. Collecting the sequential duplicate samples shortly after the environmental samples were collected produced almost the same variability as collecting the concurrent duplicate samples and the environmental samples simultaneously.

Filter Blanks

Filter blanks were processed for 226 of the 230 samples collected. This was accomplished by passing 100 mL of the sterile saline buffer solution through the filter prior to processing any dilutions of the environmental samples. While passing the saline buffer solution through the filter, every attempt was made to have the saline buffer solution come in contact with every surface that the environmental sample might touch; this would help to ensure that the equipment used to process the samples was clean. No *E. coli* colonies were present in any of the filter blanks processed.

Process Blanks

Process blanks were collected following the filtering of all of the dilutions for an environmental sample. Process blanks consisting of 100 mL of the saline buffer solution were filtered after every fourth environmental sample was processed; this was done to determine the adequacy of the equipment rinses following the filtering of each dilution. Eight of the 42 process blanks contained observable concentrations of *E. coli*. The maximum concentration of *E. coli* measured in the process blanks was 5 colonies per 100 mL. All of the process blanks that had observable concentrations of *E. coli* contained less than 1 percent of the concentration of *E. coli* measured in the environmental samples.

Field Blanks

Nine field blanks were processed on randomly selected days during the length of the study. Field blanks consisted of 250 mL of the sterile saline buffer solution that was poured into a sample-collection bottle before the first sample was taken each day. The field blanks were kept chilled and remained with the samples collected at all sites for that day. The field blank then was processed by passing 100 mL of the blank solution through the filter. None of the nine field blanks had observable concentrations of *E. coli*, indicating that there was no contamination resulting from transporting the samples.

Summary

The presence of *E. coli* in water is direct evidence of the presence of fecal contamination from warm-blooded animals and indicates the possible presence of pathogens. *E. coli* is one of the two preferred indicator bacteria used by the USEPA to determine the suitability of surface waters for recreational use. The water-quality standards for *E. coli* in recreational waters in Indiana include a single-sample standard and a geometric mean computed from five samples collected within a 30-day period.

Water samples collected from 46 stream sites in the Upper Wabash River Watershed from June through September 1998 were analyzed for concentrations of *E. coli*. Samples were collected at 21 sites during June and July and at 25 sites during August and September to allow computation of a five-sample geometric mean for each site. The five-sample geometric-mean concentrations ranged from 17 to 4,800 colonies per 100 mL, and concentrations for 43 sites exceeded the five-sample geometric-mean standard of 125 colonies per 100 mL. Of the 230 samples collected, 145 exceeded the single-sample standard of 235 colonies per 100 mL. Concentrations of *E. coli* at all sites ranged from an estimated 6 to 110,000 colonies per 100 mL during the study.

Of the 46 sites, 17 were at USGS streamflow-gaging stations. Based on records from these stations, 62 percent of the samples collected at these sites were collected at discharges above the median daily mean discharge. Statistically significant positive correlations between the concentrations of *E. coli* and discharge were determined for two sites where additional samples were collected, indicating increased concentrations of *E. coli* with greater discharge.

References Cited

- American Public Health Association, American Water Works Association, and Water Pollution Control Federation, 1992, Standard methods for the examination of water and wastewater (18th ed.), Parts 9221 through 9225: Washington, D.C., American Public Health Association, p. 9–1 to 9–73.
- Bordner, Robert, and Winter, J.A., 1978, Microbiological methods for monitoring the environment, water, and wastes: U.S. Environmental Protection Agency, EPA 600/8-78-017, 338 p.
- Bordner, Robert, Winter, J.A., and Scarpino, P.V., eds., 1978, Microbiological methods for monitoring the environment, water and wastes: U.S. Environmental Protection Agency, EPA 600/8-78-077, 357 p.
- Britton, L.J., and Greeson, P.E., eds., 1989, Methods for collection and analysis of aquatic biological and microbiological samples: U.S. Geological Survey Techniques of Water-Resources Investigations, book 5, chap. A4, 160 p.
- Helsel, D.R., and Hirsch, R.M., 1992, Statistical methods in water resources: Studies in Environmental Science, no. 49, New York, Elsevier Publishers, Inc., 522 p.
- Hoggatt, R.E., 1975, Drainage areas of Indiana streams: U.S. Geological Survey, Water Resources Division, 231 p.
- Myers, D.N., and Sylvester, M.A., 1997, Fecal indicator bacteria, in National field manual for the collection of water-quality data: U. S. Geological Survey Techniques of Water-Resources Investigations, book 9, chap. 7.1, p. FIB 1-38.
- Myers, D.N., and Wilde, F.D., eds., 1997, National field manual for the collection of water-quality data: U.S. Geological Survey Techniques of Water-Resources Investigations, book 9, chap. A7, 48 p.
- Oddi, M.J., ed., 1995, Indiana environmental rules—Water: Indiana Department of Environmental Management, p. 19
- Schneider, A.F., 1996, Physiography, in Lindsey, A.A., ed., Natural features of Indiana: Indianapolis, Indiana Academy of Science and Indiana State Library, p. 40–56.
- Stewart, J.A., Keeton, C.R., Hammil, L.E., Nguyen, H.T., and Majors, D.K., 1999, Water resources data, Indiana, water year 1998, U.S. Geological Survey Water-Data Report IN-98-1, 452 p.

Supplemental Data

(Tables 3, 4, 5, and 6)

Table 3. Water-quality data for selected sites in the Upper Wabash River Watershed in Indiana, June–September 1998

[--, no data; >, greater than; K, values estimated from non-ideal colony counts]

| Site number | Date | Time | Gage height (feet) | Discharge at time of sample collection (cubic feet per second) | Water temperature (degrees Celsius) | pH (Standard units) | Dissolved oxygen (milligrams per liter) | Specific conductance (microsiemens per centimeter at 25 degrees Celsius) | Turbidity (Nephelometric turbidity units) | <i>Escherichia coli</i> (colonies per 100 milliliters) |
|-----------------------------------|----------|------|--------------------|--|-------------------------------------|---------------------|---|--|---|--|
| 1 | 98-06-02 | 0933 | 80.01 | -- | 20.5 | 8.1 | 6.6 | 667 | 190 | 2,600 |
| | 98-06-09 | 1010 | 79.75 | -- | 17.0 | 8.2 | 7.8 | 654 | 83 | 220 |
| | 98-06-16 | 0930 | 85.98 | -- | 20.0 | 7.8 | -- | 546 | 250 | 6,300 |
| | 98-06-23 | 0935 | 90.67 | -- | 23.5 | 8.2 | 6.2 | 536 | 84 | 720 |
| | 98-06-30 | 0935 | 89.11 | -- | 20.5 | 7.3 | 5.6 | 347 | >1,000 | 110,000 |
| Five-sample geometric mean | | | | | | | | | | 3,100 |
| 2 | 98-06-02 | 1025 | 4.39 | 180 | 21.5 | 8.2 | 8.3 | 681 | 100 | 530 |
| | 98-06-09 | 1105 | 3.91 | 79 | 16.0 | 8.1 | 8.7 | 769 | 74 | 220 |
| | 98-06-16 | 1035 | 9.99 | 3,150 | 20.5 | 7.5 | -- | 502 | 150 | 7,000 |
| | 98-06-23 | 1035 | 4.93 | 345 | 24.5 | 8.1 | 7.6 | 585 | 86 | 780 |
| | 98-06-30 | 1035 | 7.91 | 1,700 | 23.0 | 7.4 | 4.0 | 305 | 920 | 23,000 |
| Five-sample geometric mean | | | | | | | | | | 1,700 |
| 3 | 98-06-02 | 1120 | 77.58 | -- | 22.0 | 8.3 | 9.1 | 688 | 64 | 140 |
| | 98-06-09 | 1209 | 77.33 | -- | 16.5 | 8.4 | 9.9 | 731 | 62 | 350 |
| | 98-06-16 | 1120 | 83.86 | -- | 21.0 | 7.6 | 6.8 | 501 | 120 | 700 |
| | 98-06-23 | 1120 | 77.31 | -- | 25.0 | 8.2 | 8.9 | 571 | 71 | 220 |
| | 98-06-30 | 1120 | 80.82 | -- | 23.0 | 7.5 | 4.1 | 253 | >1,000 | 23,000 |
| Five-sample geometric mean | | | | | | | | | | 700 |
| 4 | 98-06-03 | 1015 | 1.98 | 44 | 18.0 | 7.9 | 7.0 | 840 | 34 | 240 |
| | 98-06-10 | 0950 | 1.99 | 45 | 16.0 | 7.8 | 7.7 | 998 | 29 | 610 |
| | 98-06-17 | 0845 | 3.88 | 260 | 19.0 | 7.8 | 7.1 | 628 | 97 | 3,900 |
| | 98-06-24 | 1035 | 2.12 | 65 | 23.5 | 7.7 | 6.7 | 841 | 29 | 470 |
| | 98-07-01 | 1005 | 2.52 | 94 | 21.5 | 7.8 | 7.0 | 700 | 80 | 1,600 |
| Five-sample geometric mean | | | | | | | | | | 840 |

Table 3. Water-quality data for selected sites in the Upper Wabash River Watershed in Indiana, June–September 1998—Continued

| Site number | Date | Time | Gage height (feet) | Discharge at time of sample collection (cubic feet per second) | Water temperature (degrees Celsius) | pH (Standard units) | Dissolved oxygen (milligrams per liter) | Specific conductance (microsiemens per centimeter at 25 degrees Celsius) | Turbidity (Nephelometric turbidity units) | <i>Escherichia coli</i> (colonies per 100 milliliters) |
|-----------------------------------|----------|------|--------------------|--|-------------------------------------|---------------------|---|--|---|--|
| 5 | 98-06-03 | 0925 | 72.21 | -- | 19.5 | 7.9 | 7.8 | 662 | 18 | 53 |
| | 98-06-10 | 0908 | 71.87 | -- | 17.5 | 7.9 | 8.4 | 730 | 20 | 410 |
| | 98-06-17 | 0800 | 75.30 | -- | 20.0 | 7.4 | 7.4 | 465 | 130 | 4,200 |
| | 98-06-24 | 0945 | 76.31 | -- | 24.0 | 7.4 | 7.4 | 509 | 26 | 67 |
| | 98-07-01 | 0925 | 73.94 | -- | 25.5 | 7.7 | 6.5 | 532 | 96 | 580 |
| Five-sample geometric mean | | | | | | | | | | 320 |
| 6 | 98-06-03 | 1425 | 3.71 | 505 | 18.5 | 8.4 | 9.6 | 545 | 24 | K29 |
| | 98-06-10 | 1445 | 3.01 | 265 | 20.0 | 8.3 | 2.5 | 598 | 17 | 78 |
| | 98-06-17 | 1345 | 6.72 | 2,560 | 21.0 | 7.6 | 7.1 | 458 | 150 | 3,700 |
| | 98-06-24 | 1756 | 8.46 | 4,480 | 24.0 | 7.7 | 8.4 | 486 | 31 | 220 |
| | 98-07-01 | 1505 | 8.96 | 4,910 | 23.0 | 7.6 | 8.0 | 488 | 49 | 44 |
| Five-sample geometric mean | | | | | | | | | | 150 |
| 7 | 98-06-01 | 1300 | 1.92 | -- | 22.0 | 7.9 | 6.0 | 689 | 280 | 4,500 |
| | 98-06-08 | 1325 | 0.76 | -- | 17.5 | 7.8 | 7.4 | 790 | 83 | 170 |
| | 98-06-15 | 1010 | 5.51 | -- | 18.0 | 7.4 | 7.0 | 327 | >1,000 | 16,000 |
| | 98-06-22 | 1023 | 0.66 | -- | 22.0 | 7.7 | 6.4 | 858 | 79 | 6,800 |
| | 98-06-29 | 1025 | 6.59 | -- | 22.5 | 7.2 | 5.4 | 136 | >1,000 | 32,000 |
| Five-sample geometric mean | | | | | | | | | | 4,800 |
| 8 | 98-06-01 | 1450 | 81.10 | -- | 20.5 | 7.7 | 6.9 | 542 | >1,000 | 5,900 |
| | 98-06-09 | 0840 | 81.42 | -- | 16.0 | 7.8 | 7.8 | 729 | 46 | 180 |
| | 98-06-16 | 0840 | 82.64 | -- | 19.0 | 7.4 | 6.5 | 424 | 290 | 7,100 |
| | 98-06-23 | 0835 | 76.69 | -- | 21.5 | 7.6 | 6.5 | 680 | 57 | 610 |
| | 98-06-30 | 0845 | 84.38 | -- | 22.0 | 7.2 | 5.3 | 224 | >1,000 | 49,000 |
| Five-sample geometric mean | | | | | | | | | | 3,000 |

Table 3. Water-quality data for selected sites in the Upper Wabash River Watershed in Indiana, June–September 1998—Continued

| Site number | Date | Time | Gage height (feet) | Discharge at time of sample collection (cubic feet per second) | Water temperature (degrees Celsius) | pH (Standard units) | Dissolved oxygen (milligrams per liter) | Specific conductance (microsiemens per centimeter at 25 degrees Celsius) | Turbidity (Nephelometric turbidity units) | <i>Escherichia coli</i> (colonies per 100 milliliters) |
|-----------------------------------|----------|------|--------------------|--|-------------------------------------|---------------------|---|--|---|--|
| 9 | 98-06-02 | 1210 | 67.15 | -- | 22.5 | 7.9 | 7.1 | 538 | 250 | 630 |
| | 98-06-09 | 1302 | 66.25 | -- | 16.5 | 8.2 | 9.2 | 702 | 48 | 310 |
| | 98-06-16 | 1215 | 74.72 | -- | 19.5 | 7.6 | 8.0 | 255 | 1,140 | 19,000 |
| | 98-06-23 | 1244 | 72.45 | -- | 23.5 | 7.8 | 7.2 | 582 | 58 | 700 |
| | 98-06-30 | 1220 | 70.09 | -- | 25.0 | 8.0 | 7.0 | 740 | 280 | 4,000 |
| Five-sample geometric mean | | | | | | | | | | 1,600 |
| 10 | 98-06-04 | 1210 | 4.00 | 1,560 | 18.5 | 8.2 | 10.2 | 519 | 19 | 20 |
| | 98-06-11 | 1030 | 3.12 | 816 | 18.5 | 7.9 | 8.0 | 532 | 24 | 6,700 |
| | 98-06-18 | 0855 | 6.54 | 4,600 | 21.0 | 7.6 | 7.0 | 472 | 73 | 830 |
| | 98-06-25 | 1005 | 8.31 | 7,360 | 22.0 | 7.6 | 8.3 | 420 | 29 | 190 |
| | 98-07-02 | 0950 | 8.30 | 7,300 | 23.5 | 7.7 | 7.4 | 446 | 59 | 460 |
| Five-sample geometric mean | | | | | | | | | | 400 |
| 11 | 98-06-04 | 1255 | 2.45 | 37 | 16.5 | 8.2 | 8.7 | 704 | 6 | 70 |
| | 98-06-11 | 1110 | 2.59 | 44 | 17.0 | 8.0 | 7.8 | 657 | 18 | 2,300 |
| | 98-06-18 | 0930 | 5.21 | 490 | 18.5 | 7.7 | 7.8 | 512 | 94 | 410 |
| | 98-06-25 | 1100 | 3.15 | 105 | 22.5 | 7.9 | 7.9 | 652 | 16 | 800 |
| | 98-07-02 | 1030 | 3.25 | 120 | 19.5 | 8.0 | 8.3 | 572 | 36 | 1,600 |
| Five-sample geometric mean | | | | | | | | | | 610 |
| 12 | 98-06-01 | 1155 | 3.04 | 70 | 18.5 | 8.0 | 7.4 | 633 | 110 | 2,600 |
| | 98-06-08 | 1425 | 2.84 | 25 | 15.5 | 8.2 | 10.0 | 723 | 7 | 2,000 |
| | 98-06-15 | 1055 | 9.01 | 1,530 | 17.5 | 7.4 | 6.9 | 348 | 570 | 8,200 |
| | 98-06-22 | 1130 | 3.40 | 78 | 20.5 | 7.9 | 7.8 | 657 | 31 | 720 |
| | 98-06-29 | 1120 | 2.89 | 30 | 23.5 | 8.1 | 7.7 | 728 | 7 | 340 |
| Five-sample geometric mean | | | | | | | | | | 1,600 |

Table 3. Water-quality data for selected sites in the Upper Wabash River Watershed in Indiana, June–September 1998—Continued

| Site number | Date | Time | Gage height (feet) | Discharge at time of sample collection (cubic feet per second) | Water temperature (degrees Celsius) | pH (Standard units) | Dissolved oxygen (milligrams per liter) | Specific conductance (microsiemens per centimeter at 25 degrees Celsius) | Turbidity (Nephelometric turbidity units) | <i>Escherichia coli</i> (colonies per 100 milliliters) |
|-----------------------------------|----------|------|--------------------|--|-------------------------------------|---------------------|---|--|---|--|
| 13 | 98-06-01 | 1550 | 78.80 | -- | 22.0 | 7.8 | 6.4 | 468 | >1,000 | 5,800 |
| | 98-06-08 | 1550 | 77.68 | -- | 17.5 | 8.2 | 9.2 | 669 | 52 | 150 |
| | 98-06-15 | 1200 | 81.92 | -- | 18.5 | 7.6 | 7.1 | 344 | 850 | 9,700 |
| | 98-06-22 | 1235 | 78.36 | -- | 22.0 | 8.0 | 8.1 | 613 | 59 | 320 |
| | 98-06-29 | 1215 | 77.45 | -- | 26.5 | 8.4 | 9.0 | 685 | 18 | 460 |
| Five-sample geometric mean | | | | | | | | | | 1,000 |
| 14 | 98-06-01 | 1642 | 70.71 | -- | 21.5 | 7.7 | 6.5 | 405 | >1,000 | 4,200 |
| | 98-06-08 | 1625 | 68.98 | -- | 17.5 | 8.1 | 8.7 | 672 | 63 | 150 |
| | 98-06-15 | 1240 | 75.74 | -- | 18.5 | 7.6 | 7.1 | 367 | 520 | 8,300 |
| | 98-06-22 | 1348 | 70.19 | -- | 23.0 | 7.9 | 7.7 | 595 | 64 | 460 |
| | 98-06-29 | 1255 | 67.96 | -- | 26.5 | 8.2 | 7.9 | 687 | 17 | 130 |
| Five-sample geometric mean | | | | | | | | | | 790 |
| 15 | 98-06-02 | 1420 | 2.39 | 617 | 21.5 | 7.8 | 7.2 | 448 | 520 | 870 |
| | 98-06-08 | 1755 | 1.56 | 221 | 17.0 | 8.1 | 8.6 | 692 | 83 | 310 |
| | 98-06-15 | 1440 | 6.50 | 4,330 | 19.5 | 7.6 | 7.1 | 381 | 450 | 11,000 |
| | 98-06-22 | 1550 | 2.47 | 645 | 22.5 | 7.8 | 7.6 | 545 | 68 | 460 |
| | 98-06-29 | 1540 | 1.58 | 232 | 28.5 | 8.4 | 11.7 | 642 | 29 | 310 |
| Five-sample geometric mean | | | | | | | | | | 840 |
| 16 | 98-06-02 | 1340 | 58.99 | -- | 22.5 | 7.9 | 7.2 | 508 | 330 | 700 |
| | 98-06-09 | 1400 | 58.16 | -- | 16.5 | 8.1 | 8.9 | 734 | 69 | 500 |
| | 98-06-16 | 1310 | 65.48 | -- | 19.5 | 7.7 | 8.1 | 317 | 650 | 18,000 |
| | 98-06-23 | 1335 | 64.15 | -- | 23.0 | 7.8 | 7.5 | 606 | 59 | 430 |
| | 98-06-30 | 1310 | 58.68 | -- | 25.5 | 8.3 | 8.8 | 642 | 68 | 5,900 |
| Five-sample geometric mean | | | | | | | | | | 1,700 |

Table 3. Water-quality data for selected sites in the Upper Wabash River Watershed in Indiana, June–September 1998—Continued

| Site number | Date | Time | Gage height (feet) | Discharge at time of sample collection (cubic feet per second) | Water temperature (degrees Celsius) | pH (Standard units) | Dissolved oxygen (milligrams per liter) | Specific conductance (microsiemens per centimeter at 25 degrees Celsius) | Turbidity (Nephelometric turbidity units) | <i>Escherichia coli</i> (colonies per 100 milliliters) |
|-----------------------------------|----------|------|--------------------|--|-------------------------------------|---------------------|---|--|---|--|
| 17 | 98-06-03 | 1058 | 77.97 | -- | 16.0 | 7.8 | 8.8 | 840 | 17 | 200 |
| | 98-06-10 | 1038 | 78.05 | -- | 16.0 | 7.7 | 8.4 | 820 | 9 | 600 |
| | 98-06-17 | 0925 | 79.73 | -- | 18.5 | 7.4 | 6.3 | 625 | 120 | 6,700 |
| | 98-06-24 | 1118 | 78.00 | -- | 23.0 | 7.7 | 9.8 | 787 | 15 | 480 |
| | 98-07-01 | 1040 | 78.55 | -- | 21.5 | 7.6 | 6.7 | 616 | 58 | 1,700 |
| Five-sample geometric mean | | | | | | | | | | 920 |
| 18 | 98-06-03 | 1205 | 0.88 | 118 | 17.5 | 8.2 | 8.4 | 786 | 13 | 280 |
| | 98-06-10 | 1138 | 0.86 | 112 | 16.0 | 8.0 | 9.0 | 761 | 10 | 660 |
| | 98-06-17 | 1015 | 3.72 | 955 | 19.0 | 7.7 | 7.4 | 508 | 230 | 13,000 |
| | 98-06-24 | 1215 | 1.18 | 214 | 23.5 | 8.0 | 8.3 | 677 | 17 | 680 |
| | 98-07-01 | 1130 | 1.33 | 258 | 21.5 | 7.9 | 8.1 | 636 | 34 | 3,100 |
| Five-sample geometric mean | | | | | | | | | | 1,400 |
| 19 | 98-06-03 | 1328 | 75.43 | -- | 18.5 | 8.2 | 8.6 | 739 | 21 | 240 |
| | 98-06-10 | 1300 | 75.40 | -- | 16.0 | 8.1 | 8.9 | 734 | 14 | 370 |
| | 98-06-17 | 1100 | 78.66 | -- | 19.0 | 7.7 | 7.3 | 455 | 460 | 17,000 |
| | 98-06-24 | 1352 | 75.79 | -- | 24.0 | 8.2 | 10.0 | 680 | 24 | 430 |
| | 98-07-01 | 1210 | 75.80 | -- | 22.0 | 8.2 | 8.8 | 708 | 18 | 420 |
| Five-sample geometric mean | | | | | | | | | | 770 |
| 20 | 98-06-04 | 1000 | 76.01 | -- | 17.0 | 8.3 | 9.2 | 725 | 25 | 470 |
| | 98-06-11 | 0805 | 75.90 | -- | 17.5 | 8.1 | 8.2 | 705 | 20 | 550 |
| | 98-06-17 | 1140 | 80.28 | -- | 19.5 | 7.8 | 7.0 | 472 | 330 | 8,700 |
| | 98-06-25 | 0825 | 76.27 | -- | 24.0 | 8.2 | 8.1 | 677 | 19 | 250 |
| | 98-07-01 | 1250 | 76.47 | -- | 23.0 | 8.3 | 10.0 | 677 | 19 | 48 |
| Five-sample geometric mean | | | | | | | | | | 490 |

Table 3. Water-quality data for selected sites in the Upper Wabash River Watershed in Indiana, June–September 1998—Continued

| Site number | Date | Time | Gage height (feet) | Discharge at time of sample collection (cubic feet per second) | Water temperature (degrees Celsius) | pH (Standard units) | Dissolved oxygen (milligrams per liter) | Specific conductance (microsiemens per centimeter at 25 degrees Celsius) | Turbidity (Nephelometric turbidity units) | <i>Escherichia coli</i> (colonies per 100 milliliters) |
|-----------------------------------|----------|------|--------------------|--|-------------------------------------|---------------------|---|--|---|--|
| 21 | 98-06-04 | 1051 | 3.47 | 393 | 17.5 | 8.4 | 10.8 | 700 | 29 | 240 |
| | 98-06-11 | 0910 | 3.51 | 408 | 17.5 | 8.2 | 8.4 | 676 | 24 | 1,600 |
| | 98-06-18 | 0800 | 5.26 | 1,690 | 20.0 | 7.8 | 7.2 | 484 | 230 | 2,800 |
| | 98-06-25 | 0915 | 3.75 | 526 | 24.5 | 8.4 | 9.4 | 655 | 22 | 420 |
| | 98-07-02 | 0850 | 3.71 | 500 | 21.0 | 8.3 | 9.0 | 665 | 22 | 930 |
| Five-sample geometric mean | | | | | | | | | | 840 |
| 22 | 98-08-06 | 0910 | 82.18 | -- | 23.0 | 7.5 | 5.6 | 450 | 3 | 160 |
| | 98-08-13 | 0830 | 82.11 | -- | 24.0 | 7.4 | 6.2 | 454 | 2 | 57 |
| | 98-08-20 | 0945 | 82.38 | -- | 23.0 | 7.3 | 5.1 | 555 | 1 | K8 |
| | 98-08-27 | 0815 | 82.11 | -- | 23.5 | 7.3 | 5.3 | 481 | 2 | 28 |
| | 98-09-03 | 0935 | 82.30 | -- | 20.5 | 7.4 | 5.4 | 491 | 2 | 40 |
| Five-sample geometric mean | | | | | | | | | | 38 |
| 23 | 98-08-06 | 0955 | 6.09 | 115 | 25.5 | 8.1 | 7.1 | 389 | 3 | 130 |
| | 98-08-13 | 0900 | 6.70 | 187 | 25.5 | 8.2 | 8.1 | 385 | 2 | K8 |
| | 98-08-20 | 1030 | 5.54 | 62 | 25.5 | 8.2 | 8.6 | 454 | 2 | K6 |
| | 98-08-27 | 0845 | 5.54 | 62 | 26.0 | 7.9 | 7.9 | 379 | 3 | K13 |
| | 98-09-03 | 1025 | 5.32 | 41 | 23.5 | 7.9 | 7.2 | 387 | 1 | 16 |
| Five-sample geometric mean | | | | | | | | | | 17 |
| 24 | 98-08-06 | 1045 | 86.93 | -- | 23.0 | 7.5 | 5.2 | 478 | 11 | 1,300 |
| | 98-08-13 | 0950 | 87.00 | -- | 22.0 | 7.6 | 5.4 | 481 | 7 | 170 |
| | 98-08-20 | 1155 | 85.61 | -- | 21.5 | 7.6 | 6.3 | 641 | 8 | 220 |
| | 98-08-27 | 0950 | 85.50 | -- | 22.0 | 7.5 | 5.9 | 536 | 9 | 270 |
| | 98-09-03 | 1140 | 84.82 | -- | 19.0 | 7.7 | 7.5 | 601 | 4 | 220 |
| Five-sample geometric mean | | | | | | | | | | 310 |

Table 3. Water-quality data for selected sites in the Upper Wabash River Watershed in Indiana, June–September 1998—Continued

| Site number | Date | Time | Gage height (feet) | Discharge at time of sample collection (cubic feet per second) | Water temperature (degrees Celsius) | pH (Standard units) | Dissolved oxygen (milligrams per liter) | Specific conductance (microsiemens per centimeter at 25 degrees Celsius) | Turbidity (Nephelometric turbidity units) | <i>Escherichia coli</i> (colonies per 100 milliliters) |
|-----------------------------------|----------|------|--------------------|--|-------------------------------------|---------------------|---|--|---|--|
| 25 | 98-08-06 | 1125 | 86.29 | -- | 22.0 | 7.5 | 6.0 | 396 | 15 | 760 |
| | 98-08-13 | 1030 | 85.63 | -- | 21.5 | 7.7 | 6.3 | 478 | 5 | 110 |
| | 98-08-20 | 1245 | 83.84 | -- | 22.5 | 7.8 | 7.0 | 647 | 13 | 170 |
| | 98-08-27 | 1035 | 83.45 | -- | 22.5 | 7.8 | 7.0 | 562 | 13 | 110 |
| | 98-09-03 | 1245 | 82.89 | -- | 20.0 | 7.9 | 8.3 | 604 | 8 | 330 |
| Five-sample geometric mean | | | | | | | | | | 220 |
| 26 | 98-08-06 | 1210 | 80.90 | -- | 21.5 | 7.6 | 6.6 | 436 | 24 | 7,100 |
| | 98-08-13 | 1105 | 81.32 | -- | 21.5 | 7.8 | 6.8 | 483 | 9 | 200 |
| | 98-08-19 | 1330 | 79.07 | -- | 22.0 | 8.0 | 8.0 | 663 | 12 | 270 |
| | 98-08-27 | 1120 | 78.67 | -- | 22.0 | 7.8 | 7.6 | 581 | 11 | 250 |
| | 98-09-03 | 1320 | 78.33 | -- | 20.0 | 8.0 | 8.9 | 597 | 7 | 210 |
| Five-sample geometric mean | | | | | | | | | | 460 |
| 27 | 98-08-05 | 1405 | 83.33 | -- | 21.5 | 7.5 | 6.5 | 413 | 37 | 17,000 |
| | 98-08-12 | 1340 | 83.49 | -- | 22.5 | 7.9 | 7.9 | 476 | 14 | 250 |
| | 98-08-20 | 1520 | 81.70 | -- | 23.0 | 8.0 | 8.0 | 656 | 11 | 300 |
| | 98-08-26 | 1315 | 81.20 | -- | 23.0 | 8.0 | 8.3 | 580 | 10 | 260 |
| | 98-09-02 | 1500 | 81.03 | -- | 20.5 | 8.1 | 8.9 | 606 | 7 | 210 |
| Five-sample geometric mean | | | | | | | | | | 590 |
| 28 | 98-08-05 | 1320 | 9.36 | 1,710 | 21.5 | 7.6 | 6.6 | 440 | 34 | 7,200 |
| | 98-08-12 | 1300 | 10.51 | 2,200 | 23.0 | 7.8 | 7.5 | 465 | 14 | 170 |
| | 98-08-19 | 1425 | 7.43 | 967 | 23.0 | 8.0 | 7.8 | 652 | 15 | 220 |
| | 98-08-26 | 1230 | 6.59 | 688 | 23.0 | 8.0 | 7.9 | 574 | 11 | 190 |
| | 98-09-02 | 1415 | 6.11 | 526 | 21.0 | 8.1 | 8.6 | 598 | 7 | 140 |
| Five-sample geometric mean | | | | | | | | | | 370 |

Table 3. Water-quality data for selected sites in the Upper Wabash River Watershed in Indiana, June–September 1998—Continued

| Site number | Date | Time | Gage height (feet) | Discharge at time of sample collection (cubic feet per second) | Water temperature (degrees Celsius) | pH (Standard units) | Dissolved oxygen (milligrams per liter) | Specific conductance (microsiemens per centimeter at 25 degrees Celsius) | Turbidity (Nephelometric turbidity units) | <i>Escherichia coli</i> (colonies per 100 milliliters) |
|-----------------------------------|----------|------|--------------------|--|-------------------------------------|---------------------|---|--|---|--|
| 29 | 98-08-05 | 1145 | 81.22 | -- | 21.5 | 7.7 | 6.9 | 473 | 33 | 26,000 |
| | 98-08-12 | 1130 | 81.95 | -- | 23.0 | 7.8 | 7.9 | 455 | 16 | 260 |
| | 98-08-19 | 1325 | 80.12 | -- | 23.0 | 8.0 | 8.6 | 651 | 12 | 240 |
| | 98-08-26 | 1140 | 79.61 | -- | 23.5 | 8.1 | 8.5 | 581 | 12 | 190 |
| | 98-09-02 | 1315 | 79.32 | -- | 21.0 | 8.1 | 8.8 | 598 | 7 | 160 |
| Five-sample geometric mean | | | | | | | | | | 550 |
| 30 | 98-08-05 | 1100 | 81.80 | -- | 21.5 | 7.5 | 6.4 | 361 | 37 | 7,600 |
| | 98-08-12 | 1055 | 75.89 | -- | 22.5 | 7.9 | 7.4 | 563 | 17 | 200 |
| | 98-08-19 | 1150 | 79.06 | -- | 23.0 | 8.1 | 8.4 | 655 | 12 | 150 |
| | 98-08-26 | 1050 | 79.10 | -- | 23.5 | 8.1 | 8.3 | 579 | 10 | 300 |
| | 98-09-02 | 1145 | 78.64 | -- | 20.5 | 8.1 | 8.5 | 598 | 7 | 200 |
| Five-sample geometric mean | | | | | | | | | | 420 |
| 31 | 98-08-05 | 1015 | 77.77 | -- | 21.0 | 7.1 | 4.9 | 254 | 49 | 19,000 |
| | 98-08-12 | 1015 | 76.30 | -- | 21.0 | 7.7 | 7.3 | 576 | 21 | 360 |
| | 98-08-19 | 1100 | 75.68 | -- | 22.5 | 7.8 | 7.4 | 694 | 12 | 230 |
| | 98-08-26 | 1005 | 75.61 | -- | 22.5 | 7.7 | 6.6 | 582 | 10 | 320 |
| | 98-09-02 | 1040 | 75.61 | -- | 19.0 | 7.8 | 7.5 | 606 | 12 | 260 |
| Five-sample geometric mean | | | | | | | | | | 670 |
| 32 | 98-08-05 | 0930 | 55.86 | -- | 22.0 | 7.7 | 7.6 | 384 | 31 | 70,000 |
| | 98-08-12 | 0930 | 55.39 | -- | 23.5 | 7.8 | 7.4 | 488 | 18 | 170 |
| | 98-08-19 | 1020 | 55.36 | -- | 24.5 | 8.1 | 9.1 | 637 | 13 | 100 |
| | 98-08-26 | 0925 | 55.37 | -- | 25.0 | 7.9 | 6.8 | 567 | 13 | 54 |
| | 98-09-02 | 1005 | 55.24 | -- | 23.5 | 7.8 | 6.2 | 578 | 15 | 25 |
| Five-sample geometric mean | | | | | | | | | | 280 |

Table 3. Water-quality data for selected sites in the Upper Wabash River Watershed in Indiana, June–September 1998—Continued

| Site number | Date | Time | Gage height (feet) | Discharge at time of sample collection (cubic feet per second) | Water temperature (degrees Celsius) | pH (Standard units) | Dissolved oxygen (milligrams per liter) | Specific conductance (microsiemens per centimeter at 25 degrees Celsius) | Turbidity (Nephelometric turbidity units) | <i>Escherichia coli</i> (colonies per 100 milliliters) |
|-----------------------------------|----------|------|--------------------|--|-------------------------------------|---------------------|---|--|---|--|
| 33 | 98-08-04 | 1245 | 4.63 | 3,310 | 22.5 | 8.1 | 7.1 | 447 | 12 | 300 |
| | 98-08-11 | 1110 | 4.96 | 3,790 | 24.0 | 7.8 | 7.1 | 420 | 15 | 140 |
| | 98-08-18 | 1415 | 3.45 | 1,790 | 26.0 | 8.1 | 7.8 | 487 | 9 | 93 |
| | 98-08-25 | 1100 | 2.18 | 450 | 25.0 | 8.0 | 7 | 508 | 6 | 32 |
| | 98-09-01 | 1235 | 2.70 | 912 | 24.0 | 7.9 | 7.1 | 525 | 7 | 120 |
| Five-sample geometric mean | | | | | | | | | | 110 |
| 34 | 98-08-03 | 1445 | 79.63 | -- | 24.5 | 8.4 | 11.2 | 659 | 6 | 97 |
| | 98-08-10 | 1335 | 78.29 | -- | 23.0 | 8.0 | 8.0 | 393 | 390 | 2,700 |
| | 98-08-17 | 1508 | 79.67 | -- | 25.0 | 8.4 | 10.5 | 653 | 6 | 210 |
| | 98-08-24 | 1355 | 79.55 | -- | 26.0 | 8.2 | 11.2 | 657 | 10 | 150 |
| | 98-08-31 | 1520 | 79.46 | -- | 24.0 | 8.2 | 10.2 | 678 | 5 | 150 |
| Five-sample geometric mean | | | | | | | | | | 260 |
| 35 | 98-08-03 | 1515 | 2.10 | 91 | 25.0 | 8.4 | 11.2 | 676 | 4 | 46 |
| | 98-08-10 | 1400 | 3.16 | 314 | 23.5 | 8.1 | 7.9 | 484 | 120 | 6,400 |
| | 98-08-17 | 1540 | 2.15 | 98 | 25.5 | 8.4 | 9.7 | 656 | 7 | 100 |
| | 98-08-24 | 1430 | 1.88 | 62 | 26.5 | 8.4 | 12.5 | 663 | 6 | 64 |
| | 98-08-31 | 1543 | 1.81 | 54 | 25.0 | 8.3 | 10.4 | 685 | 8 | 67 |
| Five-sample geometric mean | | | | | | | | | | 170 |
| 36 | 98-08-04 | 1512 | 3.60 | 354 | 24.5 | 8.3 | 9.9 | 636 | 9 | 620 |
| | 98-08-10 | 1440 | 6.19 | 1,700 | 24.0 | 8.0 | 7.8 | 436 | 64 | 1,900 |
| | 98-08-17 | 1620 | 3.66 | 422 | 25.5 | 8.5 | 10.7 | 607 | 12 | 80 |
| | 98-08-24 | 1505 | 3.27 | 247 | 26.5 | 8.4 | 11.8 | 658 | 7 | 39 |
| | 98-08-31 | 1610 | 3.07 | 189 | 24.5 | 8.3 | 10.9 | 663 | 9 | 70 |
| Five-sample geometric mean | | | | | | | | | | 190 |

Table 3. Water-quality data for selected sites in the Upper Wabash River Watershed in Indiana, June–September 1998—Continued

| Site number | Date | Time | Gage height (feet) | Discharge at time of sample collection (cubic feet per second) | Water temperature (degrees Celsius) | pH (Standard units) | Dissolved oxygen (milligrams per liter) | Specific conductance (microsiemens per centimeter at 25 degrees Celsius) | Turbidity (Nephelometric turbidity units) | <i>Escherichia coli</i> (colonies per 100 milliliters) |
|-----------------------------------|----------|------|--------------------|--|-------------------------------------|---------------------|---|--|---|--|
| 37 | 98-08-04 | 1440 | 70.35 | -- | 24.5 | 7.9 | 7.0 | 380 | 25 | 14,000 |
| | 98-08-11 | 1235 | 72.26 | -- | 23.5 | 7.7 | 7.1 | 350 | 40 | 520 |
| | 98-08-18 | 1625 | 69.75 | -- | 25.0 | 7.8 | 7.5 | 357 | 18 | 83 |
| | 98-08-25 | 1235 | 67.24 | -- | 24.5 | 7.8 | 7.7 | 355 | 15 | 3,300 |
| | 98-09-01 | 1515 | 67.04 | -- | 23.5 | 7.8 | 7.6 | 355 | 14 | 220 |
| Five-sample geometric mean | | | | | | | | | | 850 |
| 38 | 98-08-04 | 1325 | 66.02 | -- | 24.0 | 7.8 | 7.2 | 351 | 32 | 1,700 |
| | 98-08-11 | 1150 | 67.62 | -- | 23.0 | 7.7 | 7.4 | 312 | 48 | 810 |
| | 98-08-18 | 1510 | 65.94 | -- | 24.5 | 7.8 | 7.7 | 318 | 20 | 240 |
| | 98-08-25 | 1135 | 63.70 | -- | 24.0 | 7.8 | 7.8 | 325 | 18 | 1,400 |
| | 98-09-01 | 1355 | 63.60 | -- | 23.0 | 7.7 | 7.9 | 327 | 14 | 36 |
| Five-sample geometric mean | | | | | | | | | | 440 |
| 39 | 98-08-04 | 0945 | 78.03 | -- | 23.5 | 7.8 | 7.0 | 376 | 42 | 4,900 |
| | 98-08-11 | 0855 | 78.65 | -- | 22.5 | 7.7 | 7.5 | 307 | 39 | 1,300 |
| | 98-08-18 | 1045 | 79.05 | -- | 24.0 | 7.6 | 7.4 | 323 | 17 | 200 |
| | 98-08-25 | 0845 | 77.07 | -- | 24.0 | 7.5 | 7.4 | 325 | 15 | 3,000 |
| | 98-09-01 | 0955 | 77.10 | -- | 23.0 | 7.6 | 7.3 | 330 | 16 | 80 |
| Five-sample geometric mean | | | | | | | | | | 790 |
| 40 | 98-08-05 | 1310 | 12.71 | 5,790 | 21.0 | 7.1 | 6.2 | 128 | 73 | 80,000 |
| | 98-08-12 | 1410 | 2.69 | 165 | 23.0 | 7.8 | 7.7 | 524 | 11 | 480 |
| | 98-08-19 | 1440 | 1.82 | 30 | 23.5 | 8.0 | 9.2 | 542 | 9 | 220 |
| | 98-08-26 | 1350 | 1.64 | 15 | 23.5 | 7.9 | 7.8 | 594 | 6 | 130 |
| | 98-09-02 | 1315 | 1.54 | 8.8 | 21.0 | 8.0 | 8.6 | 614 | 4 | 100 |
| Five-sample geometric mean | | | | | | | | | | 640 |

Table 3. Water-quality data for selected sites in the Upper Wabash River Watershed in Indiana, June–September 1998—Continued

| Site number | Date | Time | Gage height (feet) | Discharge at time of sample collection (cubic feet per second) | Water temperature (degrees Celsius) | pH (Standard units) | Dissolved oxygen (milligrams per liter) | Specific conductance (microsiemens per centimeter at 25 degrees Celsius) | Turbidity (Nephelometric turbidity units) | <i>Escherichia coli</i> (colonies per 100 milliliters) |
|-----------------------------------|----------|------|--------------------|--|-------------------------------------|---------------------|---|--|---|--|
| 41 | 98-08-03 | 1150 | 76.37 | -- | 23.5 | 8.0 | 8.1 | 619 | 9 | 210 |
| | 98-08-10 | 1045 | -- | -- | 23.0 | 7.8 | 7.2 | 469 | 28 | 600 |
| | 98-08-17 | 1155 | 76.84 | -- | 23.5 | 7.6 | 6.4 | 412 | 73 | 21,000 |
| | 98-08-24 | 1105 | 76.17 | -- | 24.5 | 7.3 | 7.2 | 738 | 7 | 500 |
| | 98-08-31 | 1125 | 75.97 | -- | 22.5 | 7.8 | 6.8 | 838 | 7 | 390 |
| Five-sample geometric mean | | | | | | | | | | 880 |
| 42 | 98-08-04 | 1030 | 79.63 | -- | 22.0 | 8.0 | 7.8 | 562 | 5 | 270 |
| | 98-08-11 | 0950 | 80.31 | -- | 21.0 | 8.0 | 8.0 | 592 | 15 | 780 |
| | 98-08-18 | 1135 | 79.46 | -- | 23.5 | 8.2 | 9.9 | 648 | 3 | 220 |
| | 98-08-25 | 0930 | 79.30 | -- | 24.0 | 7.9 | 8.1 | 630 | 5 | 730 |
| | 98-09-01 | 1045 | 79.09 | -- | 20.5 | 7.8 | 7.9 | 657 | 4 | 300 |
| Five-sample geometric mean | | | | | | | | | | 400 |
| 43 | 98-08-04 | 1115 | 2.55 | 131 | 22.5 | 8.2 | 8.6 | 636 | 5 | 1,500 |
| | 98-08-11 | 1025 | 3.15 | 292 | 22.0 | 8.1 | 8.1 | 563 | 19 | 650 |
| | 98-08-18 | 1240 | 2.40 | 99 | 24.5 | 8.3 | 10.8 | 663 | 4 | 100 |
| | 98-08-25 | 1010 | 2.23 | 68 | 24.0 | 8.1 | 8.9 | 653 | 4 | 150 |
| | 98-09-01 | 1140 | 2.11 | 51 | 20.5 | 8.1 | 8.7 | 691 | 4 | 160 |
| Five-sample geometric mean | | | | | | | | | | 300 |
| 44 | 98-08-03 | 1250 | 1.95 | 217 | 23.0 | 8.1 | 9.0 | 627 | 11 | 330 |
| | 98-08-10 | 1140 | 3.59 | 1,090 | 23.0 | 7.9 | 7.6 | 447 | 45 | 600 |
| | 98-08-17 | 1250 | 2.62 | 499 | 24.0 | 8.1 | 8.2 | 603 | 21 | 520 |
| | 98-08-24 | 1210 | 1.61 | 135 | 25.0 | 8.0 | 8.4 | 716 | 9 | 150 |
| | 98-08-31 | 1230 | 1.43 | 95 | 22.5 | 8.0 | 8.3 | 720 | 9 | 120 |
| Five-sample geometric mean | | | | | | | | | | 280 |

Table 3. Water-quality data for selected sites in the Upper Wabash River Watershed in Indiana, June–September 1998—Continued

| Site number | Date | Time | Gage height (feet) | Discharge at time of sample collection (cubic feet per second) | Water temperature (degrees Celsius) | pH (Standard units) | Dissolved oxygen (milligrams per liter) | Specific conductance (microsiemens per centimeter at 25 degrees Celsius) | Turbidity (Nephelometric turbidity units) | <i>Escherichia coli</i> (colonies per 100 milliliters) |
|-----------------------------------|----------|------|--------------------|--|-------------------------------------|---------------------|---|--|---|--|
| 45 | 98-08-03 | 1340 | 86.12 | -- | 23.0 | 8.2 | 10.1 | 648 | 5 | 260 |
| | 98-08-10 | 1225 | 87.03 | -- | 22.0 | 8.1 | 7.9 | 536 | 37 | 630 |
| | 98-08-17 | 1345 | 86.35 | -- | 24.5 | 8.3 | 9.4 | 616 | 10 | 240 |
| | 98-08-24 | 1255 | 85.96 | -- | 25.0 | 8.2 | 9.3 | 633 | 7 | 280 |
| | 98-08-31 | 1316 | 85.90 | -- | 22.0 | 8.1 | 8.6 | 633 | 8 | 220 |
| Five-sample geometric mean | | | | | | | | | | 300 |
| 46 | 98-08-03 | 1410 | 82.06 | -- | 23.0 | 8.4 | 10.7 | 854 | 7 | 180 |
| | 98-08-10 | 1250 | 82.56 | -- | 22.5 | 8.1 | 7.6 | 594 | 34 | 3,500 |
| | 98-08-17 | 1415 | 82.05 | -- | 24.5 | 8.4 | 9.9 | 879 | 16 | 340 |
| | 98-08-24 | 1320 | 82.03 | -- | 25.5 | 8.3 | 8.7 | 905 | 9 | 250 |
| | 98-08-31 | 1352 | 81.90 | -- | 23.0 | 8.2 | 8.9 | 968 | 11 | 620 |
| Five-sample geometric mean | | | | | | | | | | 510 |

Table 4. Water-quality data for two sites in the Upper Wabash River Watershed in Indiana where additional samples were collected during July–September 1998
 [--, no data; >, greater than; K, values estimated from non-ideal colony counts]

| Site number | Date | Time | Gage height (feet) | Discharge at time of sample collection (cubic feet per second) | Water temperature (degrees Celsius) | pH (Standard units) | Dissolved oxygen (milligrams per liter) | Specific conductance (microsiemens per centimeter at 25 degrees Celsius) | Turbidity (Nephelometric turbidity units) | <i>Escherichia coli</i> (colonies per 100 milliliters) |
|-------------|----------|------|--------------------|--|-------------------------------------|---------------------|---|--|---|--|
| 2 | 98-06-02 | 1025 | 4.39 | 180 | 21.5 | 8.2 | 8.3 | 681 | 100 | 530 |
| | 98-06-09 | 1105 | 3.91 | 79 | 16.2 | 8.1 | 8.7 | 769 | 74 | 220 |
| | 98-06-16 | 1035 | 9.99 | 3,150 | 20.3 | 7.5 | -- | 502 | 150 | 7,000 |
| | 98-06-23 | 1035 | 4.93 | 345 | 24.4 | 8.1 | 7.6 | 585 | 86 | 780 |
| | 98-06-30 | 1035 | 7.91 | 1,700 | 22.9 | 7.4 | 4.0 | 305 | 920 | 23,000 |
| | 98-07-22 | 1015 | 9.70 | 2,820 | 22.0 | 7.2 | 5.5 | 209 | 570 | 34,000 |
| | 98-07-29 | 1012 | 6.63 | 1,020 | 24.5 | 7.3 | 3.9 | 375 | 40 | 630 |
| | 98-08-05 | 1020 | 9.83 | 2,940 | 21.0 | 7.1 | 5.2 | 159 | 220 | 57,000 |
| | 98-08-12 | 1110 | 6.53 | 954 | 24.4 | 7.4 | 5.3 | 387 | 52 | 800 |
| | 98-08-19 | 1130 | 4.32 | 164 | 24.3 | 8.1 | 8.5 | 549 | 75 | 370 |
| | 98-08-26 | 1110 | 3.99 | 93 | 23.3 | 7.8 | 6.0 | 673 | 72 | 570 |
| | 98-09-02 | 1105 | 3.83 | 65 | 21.0 | 8.0 | 7.1 | 699 | 78 | 360 |
| | 98-09-09 | 1100 | 3.66 | 43 | 18.3 | 7.8 | 5.2 | 796 | 53 | 560 |
| 15 | 98-06-02 | 1420 | 2.39 | 617 | 21.4 | 7.8 | 7.2 | 448 | 520 | 870 |
| | 98-06-08 | 1755 | 1.56 | 221 | 16.9 | 8.1 | 8.6 | 692 | 83 | 310 |
| | 98-06-15 | 1440 | 6.50 | 4,330 | 19.3 | 7.6 | 7.1 | 381 | 450 | 11,000 |
| | 98-06-22 | 1550 | 2.47 | 645 | 22.6 | 7.8 | 7.6 | 545 | 68 | 460 |
| | 98-06-29 | 1540 | 1.58 | 232 | 28.3 | 8.4 | 11.7 | 642 | 29 | 310 |
| | 98-07-22 | 1158 | 8.47 | 6,840 | 22.3 | 7.5 | 6.7 | 206 | 520 | 43,000 |
| | 98-07-29 | 1149 | 1.91 | 378 | 23.8 | 7.9 | 7.9 | 549 | 19 | 470 |
| | 98-08-05 | 1155 | 15.92 | 21,500 | 20.9 | 7.4 | 6.8 | 115 | 210 | 31,000 |
| | 98-08-12 | 1230 | 2.49 | 680 | 23.1 | 7.8 | 7.7 | 446 | 29 | 1,800 |
| | 98-08-19 | 1305 | 1.57 | 228 | 24.2 | 8.1 | 10.0 | 606 | 19 | 250 |
| | 98-08-26 | 1225 | 1.34 | 147 | 24.2 | 8.1 | 9.6 | 708 | 22 | 750 |
| | 98-09-02 | 1230 | 1.22 | 108 | 22.2 | 8.2 | 10.7 | 731 | 21 | 240 |
| | 98-09-09 | 1210 | 1.21 | 105 | 21.0 | 8.0 | 9.4 | 714 | 19 | K830 |

Table 5. Quality-assurance data associated with the Group 1 *Escherichia coli* samples collected in the Upper Wabash River Watershed in Indiana, June–September 1998
 [K, values estimated from non-ideal colony counts; --, no data]

| Site number | Date | Time | <i>Escherichia coli</i> (colonies per 100 milliliters) | Filter blank | Process blank | Field blank | Sequential duplicate sample | Natural log percent difference |
|-------------|----------|------|---|--------------|---------------|-------------|--|--------------------------------|
| | | | | | | | for <i>Escherichia coli</i> (colonies per 100 milliliters) | |
| 1 | 98-06-02 | 0933 | 2,600 | 0 | -- | -- | -- | -- |
| | 98-06-09 | 1010 | 220 | 0 | -- | -- | -- | -- |
| | 98-06-16 | 0930 | 6,300 | 0 | 0 | -- | -- | -- |
| | 98-06-23 | 0935 | 720 | 0 | -- | -- | 700 | 3 |
| | 98-06-30 | 0935 | 110,000 | 0 | 3 | -- | -- | -- |
| 2 | 98-06-02 | 1025 | 530 | 0 | -- | -- | -- | -- |
| | 98-06-09 | 1105 | 220 | 0 | -- | -- | -- | -- |
| | 98-06-16 | 1035 | 7,000 | 0 | -- | -- | -- | -- |
| | 98-06-23 | 1035 | 780 | 0 | -- | -- | -- | -- |
| | 98-06-30 | 1035 | 23,000 | 0 | -- | -- | -- | -- |
| 2 | 98-07-22 | 1015 | 34,000 | 0 | 5 | -- | -- | -- |
| | 98-07-29 | 1012 | 630 | 0 | -- | -- | -- | -- |
| | 98-08-05 | 1020 | 57,000 | 0 | -- | -- | -- | -- |
| | 98-08-12 | 1110 | 800 | 0 | -- | -- | -- | -- |
| | 98-08-19 | 1130 | 370 | 0 | 0 | -- | -- | -- |
| 2 | 98-08-26 | 1110 | 570 | 0 | -- | -- | 340 | 52 |
| | 98-09-02 | 1105 | 360 | 0 | 0 | -- | -- | -- |
| | 98-09-09 | 1100 | 560 | 0 | -- | -- | 570 | -2 |
| 3 | 98-06-02 | 1120 | 140 | 0 | -- | -- | -- | -- |
| | 98-06-09 | 1209 | 350 | 0 | -- | -- | -- | -- |
| | 98-06-16 | 1120 | 700 | 0 | -- | -- | -- | -- |
| | 98-06-23 | 1120 | 220 | 0 | -- | -- | -- | -- |
| | 98-06-30 | 1120 | 23,000 | 0 | -- | -- | 20,000 | 14 |

Table 5. Quality-assurance data associated with the Group 1 *Escherichia coli* samples collected in the Upper Wabash River Watershed in Indiana, June–September 1998—Continued

| Site number | Date | Time | <i>Escherichia coli</i> (colonies per 100 milliliters) | Filter blank | Process blank | Field blank | Sequential duplicate sample for <i>Escherichia coli</i> (colonies per 100 milliliters) | | Natural log percent difference |
|-------------|----------|------|---|--------------|---------------|-------------|---|----|--------------------------------|
| | | | | | | | | | |
| 4 | 98-06-03 | 1015 | 240 | 0 | -- | -- | -- | -- | -- |
| | 98-06-10 | 0950 | 610 | 0 | -- | -- | -- | -- | -- |
| | 98-06-17 | 0845 | 3,900 | 0 | -- | -- | -- | -- | -- |
| | 98-06-24 | 1035 | 470 | 0 | -- | -- | -- | -- | -- |
| | 98-07-01 | 1005 | 1,600 | 0 | -- | -- | -- | -- | -- |
| 5 | 98-06-03 | 0925 | 53 | 0 | -- | -- | -- | -- | -- |
| | 98-06-10 | 0908 | 410 | 0 | -- | -- | -- | -- | -- |
| | 98-06-17 | 0800 | 4,200 | 0 | 0 | -- | -- | -- | -- |
| | 98-06-24 | 0945 | 67 | 0 | -- | 0 | 67 | 0 | 0 |
| | 98-07-01 | 0925 | 580 | 0 | -- | -- | -- | -- | -- |
| 6 | 98-06-03 | 1425 | K29 | 0 | -- | -- | -- | -- | -- |
| | 98-06-10 | 1445 | 78 | 0 | 0 | -- | 48 | 49 | 49 |
| | 98-06-17 | 1345 | 3,700 | 0 | -- | -- | -- | -- | -- |
| | 98-06-24 | 1756 | 220 | 0 | -- | -- | -- | -- | -- |
| | 98-07-01 | 1505 | 44 | 0 | -- | -- | -- | -- | -- |
| 7 | 98-06-01 | 1300 | 4,500 | 0 | -- | -- | -- | -- | -- |
| | 98-06-08 | 1325 | 170 | 0 | -- | -- | -- | -- | -- |
| | 98-06-15 | 1010 | 16,000 | -- | -- | -- | -- | -- | -- |
| | 98-06-22 | 1023 | 6,800 | 0 | -- | -- | 7,400 | -8 | -8 |
| | 98-06-29 | 1025 | 32,000 | 0 | -- | -- | -- | -- | -- |
| 8 | 98-06-01 | 1450 | 5,900 | 0 | -- | -- | -- | -- | -- |
| | 98-06-09 | 0840 | 180 | 0 | -- | -- | 130 | 33 | 33 |
| | 98-06-16 | 0840 | 7,100 | 0 | -- | -- | -- | -- | -- |
| | 98-06-23 | 0835 | 610 | 0 | -- | -- | -- | -- | -- |
| | 98-06-30 | 0845 | 49,000 | 0 | -- | -- | -- | -- | -- |

Table 5. Quality-assurance data associated with the Group 1 *Escherichia coli* samples collected in the Upper Wabash River Watershed in Indiana, June–September 1998—Continued

| Site number | Date | Time | <i>Escherichia coli</i> (colonies per 100 milliliters) | Filter blank | Process blank | Field blank | Sequential duplicate sample for <i>Escherichia coli</i> (colonies per 100 milliliters) | Natural log percent difference |
|-------------|----------|------|---|--------------|---------------|-------------|---|--------------------------------|
| 9 | 98-06-02 | 1210 | 630 | 0 | -- | -- | -- | -- |
| | 98-06-09 | 1302 | 310 | 0 | 0 | -- | -- | -- |
| | 98-06-16 | 1215 | 19,000 | 0 | -- | -- | 20,000 | -5 |
| | 98-06-23 | 1244 | 700 | 0 | 0 | -- | -- | -- |
| | 98-06-30 | 1220 | 4,000 | 0 | -- | -- | -- | -- |
| 10 | 98-06-04 | 1210 | 20 | 0 | -- | -- | -- | -- |
| | 98-06-11 | 1030 | 6,700 | 0 | -- | -- | -- | -- |
| | 98-06-18 | 0855 | 830 | 0 | 2 | -- | 870 | -5 |
| | 98-06-25 | 1005 | 190 | 0 | -- | -- | -- | -- |
| | 98-07-02 | 0950 | 460 | 0 | -- | -- | -- | -- |
| 11 | 98-06-04 | 1255 | 70 | 0 | 0 | 0 | 80 | -13 |
| | 98-06-11 | 1110 | 2,300 | 0 | -- | -- | -- | -- |
| | 98-06-18 | 0930 | 410 | 0 | 3 | -- | -- | -- |
| | 98-06-25 | 1100 | 800 | 0 | 0 | -- | -- | -- |
| | 98-07-02 | 1030 | 1,600 | 0 | -- | -- | 1,400 | 13 |
| 12 | 98-06-01 | 1155 | 2,600 | 0 | -- | -- | -- | -- |
| | 98-06-08 | 1425 | 2,000 | 0 | -- | -- | -- | -- |
| | 98-06-15 | 1055 | 8,200 | 0 | 2 | 0 | 7,500 | 9 |
| | 98-06-22 | 1130 | 720 | 0 | -- | -- | -- | -- |
| | 98-06-29 | 1120 | 340 | 0 | 0 | -- | 280 | 19 |
| 13 | 98-06-01 | 1550 | 5,800 | 0 | -- | -- | -- | -- |
| | 98-06-08 | 1550 | 150 | 0 | 0 | -- | 160 | -6 |
| | 98-06-15 | 1200 | 9,700 | 0 | -- | -- | -- | -- |
| | 98-06-22 | 1235 | 320 | 0 | -- | -- | -- | -- |
| | 98-06-29 | 1215 | 460 | 0 | -- | -- | -- | -- |

Table 5. Quality-assurance data associated with the Group 1 *Escherichia coli* samples collected in the Upper Wabash River Watershed in Indiana, June–September 1998—Continued

| Site number | Date | Time | <i>Escherichia coli</i> (colonies per 100 milliliters) | Filter blank | Process blank | Field blank | Sequential duplicate sample for <i>Escherichia coli</i> (colonies per 100 milliliters) | Natural log percent difference |
|-------------|----------|------|---|--------------|---------------|-------------|---|-----------------------------------|
| 14 | 98-06-01 | 1642 | 4,200 | 0 | -- | -- | -- | -- |
| | 98-06-08 | 1625 | 150 | 0 | -- | -- | -- | -- |
| | 98-06-15 | 1240 | 8,300 | 0 | -- | -- | -- | -- |
| | 98-06-22 | 1348 | 460 | -- | -- | -- | -- | -- |
| | 98-06-29 | 1255 | 130 | 0 | -- | 0 | -- | -- |
| 15 | 98-06-02 | 1420 | 870 | 0 | 0 | -- | 1,300 | -40 |
| | 98-06-08 | 1755 | 310 | 0 | -- | -- | -- | -- |
| | 98-06-15 | 1440 | 11,000 | 0 | -- | -- | -- | -- |
| | 98-06-22 | 1550 | 460 | 0 | 0 | -- | -- | -- |
| | 98-06-29 | 1540 | 310 | 0 | -- | -- | -- | -- |
| 15 | 98-07-22 | 1158 | 43,000 | 0 | -- | -- | -- | -- |
| | 98-07-29 | 1149 | 470 | 0 | 0 | -- | -- | -- |
| | 98-08-05 | 1155 | 31,000 | 0 | -- | -- | -- | -- |
| | 98-08-12 | 1230 | 1,800 | 0 | 0 | -- | -- | -- |
| | 98-08-19 | 1305 | 250 | 0 | -- | -- | -- | -- |
| 15 | 98-08-26 | 1225 | 750 | 0 | 0 | -- | -- | -- |
| | 98-09-02 | 1230 | 240 | 0 | -- | -- | 240 | 0 |
| | 98-09-09 | 1210 | K830 | 0 | 0 | -- | -- | -- |
| 16 | 98-06-02 | 1340 | 700 | 0 | -- | -- | -- | -- |
| | 98-06-09 | 1400 | 500 | 0 | -- | -- | -- | -- |
| | 98-06-16 | 1310 | 18,000 | 0 | -- | -- | -- | -- |
| | 98-06-23 | 1335 | 430 | 0 | -- | -- | -- | -- |
| | 98-06-30 | 1310 | 5,900 | 0 | -- | -- | -- | -- |

Table 5. Quality-assurance data associated with the Group 1 *Escherichia coli* samples collected in the Upper Wabash River Watershed in Indiana, June–September 1998—Continued

| Site number | Date | Time | <i>Escherichia coli</i> (colonies per 100 milliliters) | Filter blank | Process blank | Field blank | Sequential duplicate sample for | Natural log percent difference |
|-------------|----------|------|---|--------------|---------------|-------------|---|--------------------------------|
| | | | | | | | <i>Escherichia coli</i> (colonies per 100 milliliters) | |
| 17 | 98-06-03 | 1058 | 200 | 0 | -- | -- | -- | -- |
| | 98-06-10 | 1038 | 600 | 0 | -- | -- | -- | -- |
| | 98-06-17 | 0925 | 6,700 | 0 | -- | -- | 6,900 | -3 |
| | 98-06-24 | 1118 | 480 | 0 | -- | -- | -- | -- |
| | 98-07-01 | 1040 | 1,700 | 0 | -- | -- | 1,800 | -6 |
| 18 | 98-06-03 | 1205 | 280 | 0 | -- | -- | -- | -- |
| | 98-06-10 | 1138 | 660 | 0 | -- | -- | -- | -- |
| | 98-06-17 | 1015 | 13,000 | 0 | -- | -- | -- | -- |
| | 98-06-24 | 1215 | 680 | 0 | 0 | -- | -- | -- |
| | 98-07-01 | 1130 | 3,100 | 0 | -- | -- | -- | -- |
| 19 | 98-06-03 | 1328 | 240 | 0 | 0 | -- | 290 | -19 |
| | 98-06-10 | 1300 | 370 | 0 | -- | -- | -- | -- |
| | 98-06-17 | 1100 | 17,000 | 0 | -- | -- | -- | -- |
| | 98-06-24 | 1352 | 430 | 0 | -- | -- | -- | -- |
| | 98-07-01 | 1210 | 420 | 0 | -- | -- | -- | -- |
| 20 | 98-06-04 | 1000 | 470 | 0 | -- | -- | -- | -- |
| | 98-06-11 | 0805 | 550 | 0 | -- | -- | -- | -- |
| | 98-06-17 | 1140 | 8,700 | 0 | 0 | -- | -- | -- |
| | 98-06-25 | 0825 | 250 | 0 | -- | -- | 270 | -8 |
| | 98-07-01 | 1250 | 48 | 0 | -- | -- | -- | -- |
| 21 | 98-06-04 | 1051 | 240 | 0 | -- | -- | -- | -- |
| | 98-06-11 | 0910 | 1,600 | 0 | 4 | 0 | 800 | 69 |
| | 98-06-18 | 0800 | 2,800 | 0 | -- | -- | 3,800 | -31 |
| | 98-06-25 | 0915 | 420 | 0 | 0 | -- | -- | -- |
| | 98-07-02 | 0850 | 930 | 0 | -- | -- | -- | -- |

Table 6. Quality-assurance data associated with the Group 2 *Escherichia coli* samples collected in the Upper Wabash River Watershed in Indiana, August–September 1998

[K, values estimated from non-ideal colony counts; --, no data]

| Site number | Date | Time | <i>Escherichia coli</i> (colonies per 100 milliliters) | Filter blank | Process blank | Field blank | Concurrent duplicate sample for <i>Escherichia coli</i> (colonies per 100 milliliters) | | Natural log percent difference |
|-------------|----------|------|---|--------------|---------------|-------------|---|----|-----------------------------------|
| | | | | | | | | | |
| 22 | 98-08-06 | 0910 | 160 | 0 | -- | -- | -- | -- | -- |
| | 98-08-13 | 0830 | 57 | 0 | -- | -- | -- | -- | -- |
| | 98-08-20 | 0945 | K8 | 0 | -- | -- | K9 | -- | -12 |
| | 98-08-27 | 0815 | 28 | 0 | -- | -- | -- | -- | -- |
| | 98-09-03 | 0935 | 40 | 0 | -- | -- | -- | -- | -- |
| 23 | 98-08-06 | 0955 | 130 | 0 | -- | -- | -- | -- | -- |
| | 98-08-13 | 0900 | K8 | 0 | -- | -- | K3 | -- | 98 |
| | 98-08-20 | 1030 | K6 | 0 | -- | -- | -- | -- | -- |
| | 98-08-27 | 0845 | K13 | 0 | -- | -- | -- | -- | -- |
| | 98-09-03 | 1025 | 16 | 0 | -- | -- | -- | -- | -- |
| 24 | 98-08-06 | 1045 | 1,300 | 0 | -- | -- | -- | -- | -- |
| | 98-08-13 | 0950 | 170 | 0 | -- | -- | -- | -- | -- |
| | 98-08-20 | 1155 | 220 | 0 | -- | -- | -- | -- | -- |
| | 98-08-27 | 0950 | 270 | 0 | -- | -- | -- | -- | -- |
| | 98-09-03 | 1140 | 220 | 0 | -- | -- | 220 | -- | 0 |
| 25 | 98-08-06 | 1125 | 760 | 0 | -- | -- | -- | -- | -- |
| | 98-08-13 | 1030 | 110 | 0 | -- | -- | -- | -- | -- |
| | 98-08-20 | 1245 | 170 | 0 | -- | -- | -- | -- | -- |
| | 98-08-27 | 1035 | 110 | 0 | -- | -- | 250 | -- | -82 |
| | 98-09-03 | 1245 | 330 | 0 | -- | -- | -- | -- | -- |
| 26 | 98-08-06 | 1210 | 7,100 | 0 | -- | 0 | 2,800 | -- | 93 |
| | 98-08-13 | 1105 | 200 | 0 | -- | -- | -- | -- | -- |
| | 98-08-19 | 1330 | 270 | 0 | 0 | -- | -- | -- | -- |
| | 98-08-27 | 1120 | 250 | 0 | -- | -- | -- | -- | -- |
| | 98-09-03 | 1320 | 210 | 0 | -- | -- | -- | -- | -- |

Table 6. Quality-assurance data associated with the Group 2 *Escherichia coli* samples collected in the Upper Wabash River Watershed in Indiana, August–September 1998—Continued

| Site number | Date | Time | <i>Escherichia coli</i> (colonies per 100 milliliters) | Filter blank | Process blank | Field blank | Concurrent duplicate sample | Natural log percent difference |
|-------------|----------|------|---|--------------|---------------|-------------|--|--------------------------------|
| | | | | | | | for <i>Escherichia coli</i> (colonies per 100 milliliters) | |
| 27 | 98-08-05 | 1405 | 17,000 | 0 | -- | -- | 18,000 | -6 |
| | 98-08-12 | 1340 | 250 | 0 | -- | -- | -- | -- |
| | 98-08-20 | 1520 | 300 | 0 | 0 | -- | -- | -- |
| | 98-08-26 | 1315 | 260 | 0 | -- | -- | -- | -- |
| | 98-09-02 | 1500 | 210 | 0 | 0 | -- | -- | -- |
| 28 | 98-08-05 | 1320 | 7,200 | 0 | -- | -- | -- | -- |
| | 98-08-12 | 1300 | 170 | 0 | -- | -- | -- | -- |
| | 98-08-19 | 1425 | 220 | 0 | -- | 0 | 210 | 5 |
| | 98-08-26 | 1230 | 190 | 0 | -- | -- | -- | -- |
| | 98-09-02 | 1415 | 140 | 0 | -- | -- | -- | -- |
| 29 | 98-08-05 | 1145 | 26,000 | 0 | -- | -- | -- | -- |
| | 98-08-12 | 1130 | 260 | 0 | -- | -- | -- | -- |
| | 98-08-19 | 1325 | 240 | 0 | -- | -- | -- | -- |
| | 98-08-26 | 1140 | 190 | 0 | -- | -- | -- | -- |
| | 98-09-02 | 1315 | 160 | 0 | -- | -- | -- | -- |
| 30 | 98-08-05 | 1100 | 7,600 | 0 | -- | -- | -- | -- |
| | 98-08-12 | 1055 | 200 | 0 | -- | -- | -- | -- |
| | 98-08-19 | 1150 | 150 | 0 | -- | -- | -- | -- |
| | 98-08-26 | 1050 | 300 | 0 | -- | -- | -- | -- |
| | 98-09-02 | 1145 | 200 | 0 | -- | -- | 200 | 0 |
| 31 | 98-08-05 | 1015 | 19,000 | 0 | -- | -- | -- | -- |
| | 98-08-12 | 1015 | 360 | 0 | -- | -- | -- | -- |
| | 98-08-19 | 1100 | 230 | 0 | -- | -- | -- | -- |
| | 98-08-26 | 1005 | 320 | 0 | -- | -- | 320 | 0 |
| | 98-09-02 | 1040 | 260 | 0 | -- | -- | -- | -- |

Table 6. Quality-assurance data associated with the Group 2 *Escherichia coli* samples collected in the Upper Wabash River Watershed in Indiana, August–September 1998—Continued

| Site number | Date | Time | <i>Escherichia coli</i> (colonies per 100 milliliters) | Filter blank | Process blank | Field blank | Concurrent duplicate sample for <i>Escherichia coli</i> (colonies per 100 milliliters) | | Natural log percent difference |
|-------------|----------|------|---|--------------|---------------|-------------|---|----|-----------------------------------|
| | | | | | | | | | |
| 32 | 98-08-05 | 0930 | 70,000 | 0 | -- | -- | -- | -- | -- |
| | 98-08-12 | 0930 | 170 | 0 | -- | 0 | 170 | 0 | 0 |
| | 98-08-19 | 1020 | 100 | 0 | -- | -- | -- | -- | -- |
| | 98-08-26 | 0925 | 54 | 0 | 0 | -- | -- | -- | -- |
| | 98-09-02 | 1005 | 25 | 0 | -- | -- | -- | -- | -- |
| 33 | 98-08-04 | 1245 | 300 | 0 | -- | -- | -- | -- | -- |
| | 98-08-11 | 1110 | 140 | 0 | -- | -- | -- | -- | -- |
| | 98-08-18 | 1415 | 93 | 0 | -- | -- | -- | -- | -- |
| | 98-08-25 | 1100 | 32 | 0 | -- | -- | -- | -- | -- |
| | 98-09-01 | 1235 | 120 | 0 | -- | -- | -- | -- | -- |
| 34 | 98-08-03 | 1445 | 97 | 0 | -- | -- | -- | -- | -- |
| | 98-08-10 | 1335 | 2,700 | 0 | -- | -- | -- | -- | -- |
| | 98-08-17 | 1508 | 210 | 0 | -- | -- | 180 | 15 | 15 |
| | 98-08-24 | 1355 | 150 | 0 | -- | -- | -- | -- | -- |
| | 98-08-31 | 1520 | 150 | 0 | -- | -- | -- | -- | -- |
| 35 | 98-08-03 | 1515 | 46 | 0 | -- | -- | 31 | 40 | 40 |
| | 98-08-10 | 1400 | 6,400 | 0 | -- | -- | -- | -- | -- |
| | 98-08-17 | 1540 | 100 | 0 | -- | -- | -- | -- | -- |
| | 98-08-24 | 1430 | 64 | 0 | -- | -- | -- | -- | -- |
| | 98-08-31 | 1543 | 67 | 0 | -- | -- | -- | -- | -- |
| 36 | 98-08-04 | 1512 | 620 | 0 | -- | -- | 280 | 80 | 80 |
| | 98-08-10 | 1440 | 1,900 | 0 | -- | -- | -- | -- | -- |
| | 98-08-17 | 1620 | 80 | 0 | 0 | -- | -- | -- | -- |
| | 98-08-24 | 1505 | 39 | 0 | -- | -- | -- | -- | -- |
| | 98-08-31 | 1610 | 70 | 0 | 0 | -- | -- | -- | -- |

Table 6. Quality-assurance data associated with the Group 2 *Escherichia coli* samples collected in the Upper Wabash River Watershed in Indiana, August–September 1998—Continued

| Site number | Date | Time | <i>Escherichia coli</i> (colonies per 100 milliliters) | Filter blank | Process blank | Field blank | Concurrent duplicate sample for <i>Escherichia coli</i> (colonies per 100 milliliters) | | Natural log percent difference |
|-------------|----------|------|---|--------------|---------------|-------------|---|----|--------------------------------|
| | | | | | | | | | |
| 37 | 98-08-04 | 1440 | 14,000 | 0 | -- | -- | -- | -- | -- |
| | 98-08-11 | 1235 | 520 | 0 | -- | -- | -- | -- | -- |
| | 98-08-18 | 1625 | 83 | 0 | 0 | -- | 90 | -- | -8 |
| | 98-08-25 | 1235 | 3,300 | 0 | -- | -- | -- | -- | -- |
| | 98-09-01 | 1515 | 220 | 0 | -- | -- | -- | -- | -- |
| 38 | 98-08-04 | 1325 | 1,700 | 0 | -- | -- | -- | -- | -- |
| | 98-08-11 | 1150 | 810 | 0 | -- | -- | -- | -- | -- |
| | 98-08-18 | 1510 | 240 | 0 | -- | -- | -- | -- | -- |
| | 98-08-25 | 1135 | 1,400 | 0 | -- | -- | -- | -- | -- |
| | 98-09-01 | 1355 | 36 | 0 | 0 | 0 | 33 | -- | 9 |
| 39 | 98-08-04 | 0945 | 4,900 | 0 | -- | -- | -- | -- | -- |
| | 98-08-11 | 0855 | 1,300 | 0 | -- | -- | 300 | -- | 147 |
| | 98-08-18 | 1045 | 200 | 0 | -- | -- | -- | -- | -- |
| | 98-08-25 | 0845 | 3,000 | 0 | -- | -- | -- | -- | -- |
| | 98-09-01 | 0955 | 80 | 0 | -- | -- | -- | -- | -- |
| 40 | 98-08-05 | 1310 | 80,000 | 0 | 0 | -- | -- | -- | -- |
| | 98-08-12 | 1410 | 480 | 0 | -- | -- | -- | -- | -- |
| | 98-08-19 | 1440 | 220 | 0 | -- | -- | 200 | -- | 10 |
| | 98-08-26 | 1350 | 130 | 0 | -- | -- | -- | -- | -- |
| | 98-09-02 | 1315 | 100 | 0 | 0 | -- | -- | -- | -- |
| 41 | 98-08-03 | 1150 | 210 | 0 | -- | -- | -- | -- | -- |
| | 98-08-10 | 1045 | 600 | 0 | -- | -- | 1,300 | -- | -77 |
| | 98-08-17 | 1155 | 21,000 | 0 | -- | -- | -- | -- | -- |
| | 98-08-24 | 1105 | 500 | 0 | 2 | -- | -- | -- | -- |
| | 98-08-31 | 1125 | 390 | 0 | -- | -- | -- | -- | -- |

Table 6. Quality-assurance data associated with the Group 2 *Escherichia coli* samples collected in the Upper Wabash River Watershed in Indiana, August–September 1998—Continued

| Site number | Date | Time | <i>Escherichia coli</i> (colonies per 100 milliliters) | Filter blank | Process blank | Field blank | Concurrent duplicate sample for <i>Escherichia coli</i> (colonies per 100 milliliters) | | Natural log percent difference |
|-------------|----------|------|---|--------------|---------------|-------------|---|----|-----------------------------------|
| | | | | | | | | | |
| 42 | 98-08-04 | 1030 | 270 | 0 | -- | -- | -- | -- | -- |
| | 98-08-11 | 0950 | 780 | 0 | -- | -- | -- | -- | -- |
| | 98-08-18 | 1135 | 220 | 0 | -- | -- | -- | -- | -- |
| | 98-08-25 | 0930 | 730 | 0 | 0 | -- | -- | -- | -- |
| | 98-09-01 | 1045 | 300 | 0 | -- | -- | -- | -- | -- |
| 43 | 98-08-04 | 1115 | 1,500 | 0 | 0 | -- | -- | -- | -- |
| | 98-08-11 | 1025 | 650 | 0 | 1 | -- | -- | -- | -- |
| | 98-08-18 | 1240 | 100 | 0 | -- | -- | -- | -- | -- |
| | 98-08-25 | 1010 | 150 | 0 | -- | -- | 93 | 48 | 48 |
| | 98-09-01 | 1140 | 160 | 0 | -- | -- | -- | -- | -- |
| 44 | 98-08-03 | 1250 | 330 | 0 | -- | -- | -- | -- | -- |
| | 98-08-10 | 1140 | 600 | 0 | -- | -- | -- | -- | -- |
| | 98-08-17 | 1250 | 520 | 0 | -- | -- | -- | -- | -- |
| | 98-08-24 | 1210 | 150 | 0 | -- | -- | 150 | 0 | 0 |
| | 98-08-31 | 1230 | 120 | 0 | -- | -- | -- | -- | - |
| 45 | 98-08-03 | 1340 | 260 | 0 | 0 | -- | -- | -- | -- |
| | 98-08-10 | 1225 | 630 | 0 | -- | -- | -- | -- | -- |
| | 98-08-17 | 1345 | 240 | 0 | -- | -- | -- | -- | -- |
| | 98-08-24 | 1255 | 280 | 0 | -- | -- | -- | -- | -- |
| | 98-08-31 | 1316 | 220 | 0 | -- | -- | 173 | 24 | 24 |
| 46 | 98-08-03 | 1410 | 180 | 0 | -- | -- | -- | -- | -- |
| | 98-08-10 | 1250 | 3,500 | 0 | -- | -- | -- | -- | -- |
| | 98-08-17 | 1415 | 340 | 0 | -- | -- | -- | -- | -- |
| | 98-08-24 | 1320 | 250 | 0 | -- | -- | -- | -- | -- |
| | 98-08-31 | 1352 | 620 | 0 | -- | -- | -- | -- | -- |